


GPx-5 (D-3): sc-390092



The Power to Question

BACKGROUND

Glutathione peroxidase (GPx) enzymes are generally selenium-containing tetrameric glycoproteins that help prevent lipid peroxidation of cell membranes. GPx enzymes reduce lipid hydroperoxides to alcohols, and reduce free hydrogen peroxide to water. GPx members are among the few proteins known in higher vertebrates to contain selenocysteine, which occurs at the active site of glutathione peroxidase and is coded by the nonsense (stop) codon TGA. There are eight GPx homologs (GPx-1-8). GPx-1 plays an important role in the antioxidant defense of the vascular wall and neural cells in response to oxidative stress. GPx-2 is the major isoform in the lungs and its basal or inducible expression is dependent on Nrf2. GPx-3 is under regulation by hypoxic stress and the expression and deficiency of GPx-3 is associated with cardiovascular disease and stroke. GPx-5 is selenium-independent; it is bound to the acrosome of sperm, where it may protect sperm from premature acrosome reaction in the epididymis.

REFERENCES

1. Chu, F.F., et al. 1997. Expression and chromosomal mapping of mouse Gpx2 gene encoding the gastrointestinal form of glutathione peroxidase, GPX-GI. *Biomed. Environ. Sci.* 10: 156-162.
2. Hall, L., et al. 1998. The majority of human glutathione peroxidase type 5 (GPX5) transcripts are incorrectly spliced: implications for the role of GPX5 in the male reproductive tract. *Biochem. J.* 333: 5-9.
3. Bilodeau, J.F., et al. 1999. Increased resistance of GPx-1 transgenic mice to tumor promoter-induced loss of glutathione peroxidase activity in skin. *Int. J. Cancer* 80: 863-867.
4. Mork, H., et al. 2000. Inverse mRNA expression of the selenocysteine-containing proteins GI-GPx and SeP in colorectal adenomas compared with adjacent normal mucosa. *Nutr. Cancer* 37: 108-116.

CHROMOSOMAL LOCATION

Genetic locus: GPX5 (human) mapping to 6p22.1; Gpx5 (mouse) mapping to 13 A3.1.

SOURCE

GPx-5 (D-3) is a mouse monoclonal antibody raised against amino acids 22-66 mapping near the N-terminus of GPx-5 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

GPx-5 (D-3) is available conjugated to agarose (sc-390092 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390092 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390092 PE), fluorescein (sc-390092 FITC), Alexa Fluor® 488 (sc-390092 AF488), Alexa Fluor® 546 (sc-390092 AF546), Alexa Fluor® 594 (sc-390092 AF594) or Alexa Fluor® 647 (sc-390092 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-390092 AF680) or Alexa Fluor® 790 (sc-390092 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

GPx-5 (D-3) is recommended for detection of GPx-5 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GPx-5 siRNA (h): sc-62419, GPx-5 siRNA (m): sc-62420, GPx-5 shRNA Plasmid (h): sc-62419-SH, GPx-5 shRNA Plasmid (m): sc-62420-SH, GPx-5 shRNA (h) Lentiviral Particles: sc-62419-V and GPx-5 shRNA (m) Lentiviral Particles: sc-62420-V.


Molecular Weight of GPx-5: 25 kDa.

Positive Controls: mouse epididymis extract: sc-364240 or rat epididymis extract: sc-364804.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



GPx-5 (D-3): sc-390092. Western blot analysis of GPx-5 expression in mouse epididymis (A) and rat epididymis (B) tissue extracts.



GPx-5 (D-3): sc-390092. Immunoperoxidase staining of formalin fixed, paraffin-embedded human epididymis tissue showing cytoplasmic staining of glandular cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded mouse testis tissue showing cytoplasmic staining of Leydig cells (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.