SANTA CRUZ BIOTECHNOLOGY, INC.

NMDAe1 (D-8): sc-390094



BACKGROUND

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neuro-degeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for fast excitatory neurotransmission by glutamate, whereas the NMDA receptors exhibit slow kinesis of Ca²⁺ ions and a high permeability for Ca²⁺ ions. The NMDA receptors consist of five subunits: ϵ 1, 2, 3, 4 and one ζ subunit. The ζ subunit is expressed throughout the brainstem whereas the four ϵ subunits display limited distribution.

CHROMOSOMAL LOCATION

Genetic locus: GRIN2A (human) mapping to 16p13.2; Grin2a (mouse) mapping to 16 A1.

SOURCE

 $NMDA\varepsilon1$ (D-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 31-69 within an N-terminal extracellular domain of $NMDA\varepsilon1$ of human origin.

PRODUCT

Each vial contains 200 μg lgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-390094 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

NMDA ε 1 (D-8) is recommended for detection of NMDA ε 1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NMDA ϵ 1 siRNA (h): sc-36083, NMDA ϵ 1 siRNA (m): sc-36084, NMDA ϵ 1 siRNA (r): sc-270157, NMDA ϵ 1 shRNA Plasmid (h): sc-36083-SH, NMDA ϵ 1 shRNA Plasmid (m): sc-36084-SH, NMDA ϵ 1 shRNA Plasmid (r): sc-270157-SH, NMDA ϵ 1 shRNA (h) Lentiviral Particles: sc-36083-V, NMDA ϵ 1 shRNA (m) Lentiviral Particles: sc-36084-V and NMDA ϵ 1 shRNA (r) Lentiviral Particles: sc-270157-V.

Molecular Weight of NMDAE1: 177 kDa.

Positive Controls: H4 cell lysate: sc-2408, mouse brain extract: sc-2253 or rat brain extract: sc-2392.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



NINDAET (D-8): SC-390094. Western blot analysis of NMDA ϵ 1 expression in rat brain (**A**) and mouse brain (**B**) tissue extracts.

SELECT PRODUCT CITATIONS

- Brai, E., et al. 2015. Notch1 regulates hippocampal plasticity through interaction with the Reelin pathway, glutamatergic transmission and CREB signaling. Front. Cell. Neurosci. 9: 447.
- Banerjee, J., et al. 2017. Altered glutamatergic tone reveals two distinct resting state networks at the cellular level in hippocampal sclerosis. Sci. Rep. 7: 319.
- Tevzadze, G., et al. 2020. Gut neurotoxin p-cresol induces differential expression of GLUN2B and GLUN2A subunits of the NMDA receptor in the hippocampus and nucleus accumbens in healthy and audiogenic seizure-prone rats. AIMS Neurosci. 7: 30-42.
- Raymundi, A.M., et al. 2022. Effects of δ-9 tetrahydrocannabinol on fear memory labilization and reconsolidation: a putative role of GluN2B-NMDA receptor within the dorsal hippocampus. Neuropharmacology 225: 109386.
- 5. Francija, E., et al. 2022. GluN2A-ERK-mTOR pathway confers a vulnerability to LPS-induced depressive-like behaviour. Behav. Brain Res. 417: 113625.
- Glavonic, E., et al. 2023. Sex-specific role of hippocampal NMDA-ErkmTOR signaling in fear extinction of adolescent mice. Brain Res. Bull. 192: 156-167.
- 7. Alzu'bi, A., et al. 2024. Delineating the molecular mechanisms of hippocampal neurotoxicity induced by chronic administration of synthetic cannabinoid AB-FUBINACA in mice. Neurotoxicology 103: 50-59.

RESEARCH USE

For research use only, not for use in diagnostic procedures.