

PP2C ζ (C-4): sc-390214

BACKGROUND

The phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/threonine protein phosphatases. The PP2C group of serine/threonine phosphatases are divided into subclasses according to their requirement for magnesium substrate, their structure and by insensitivity to okadaic acid. PP2C ζ (protein phosphatase 2C isoform ζ), also known as PPM1J (protein phosphatase, Mg²⁺/Mn²⁺ dependent, 1J), is a 505 amino acid phosphoprotein that contains one PP2C-like domain and belongs to the PP2C family. Two isoforms of PP2C ζ are produced as a result of alternative splicing events. PP2C ζ likely exhibits its specific role through its small ubiquitin-related modifier-1-induced recruitment to UBC9 (ubiquitin conjugating enzyme 9).

REFERENCES

- Cohen, P. and Cohen, P.T. 1989. Protein phosphatases come of age. *J. Biol. Chem.* 264: 21435-21438.
- Hanada, M., et al. 1998. Selective suppression of stress-activated protein kinase pathway by protein phosphatase 2C in mammalian cells. *FEBS Lett.* 437: 172-176.
- Zolnierowicz, S. 2000. Type 2A protein phosphatase, the complex regulator of numerous signaling pathways. *Biochem. Pharmacol.* 60: 1225-1235.
- Hanada, M., et al. 2001. Regulation of the TAK1 signaling pathway by protein phosphatase 2C. *J. Biol. Chem.* 276: 5753-5759.
- Komaki, K., et al. 2003. Molecular cloning of PP2C η , a novel member of the protein phosphatase 2C family. *Biochim. Biophys. Acta* 1630: 130-137.

CHROMOSOMAL LOCATION

Genetic locus: PPM1J (human) mapping to 1p13.2; Ppm1j (mouse) mapping to 3 F2.2.

SOURCE

PP2C ζ (C-4) is a mouse monoclonal antibody raised against amino acids 431-475 mapping near the C-terminus of PP2C ζ of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

PP2C ζ (C-4) is available conjugated to agarose (sc-390214 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390214 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390214 PE), fluorescein (sc-390214 FITC), Alexa Fluor[®] 488 (sc-390214 AF488), Alexa Fluor[®] 546 (sc-390214 AF546), Alexa Fluor[®] 594 (sc-390214 AF594) or Alexa Fluor[®] 647 (sc-390214 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-390214 AF680) or Alexa Fluor[®] 790 (sc-390214 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

PP2C ζ (C-4) is recommended for detection of PP2C ζ of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PP2C ζ siRNA (h): sc-78774, PP2C ζ siRNA (m): sc-155945, PP2C ζ shRNA Plasmid (h): sc-78774-SH, PP2C ζ shRNA Plasmid (m): sc-155945-SH, PP2C ζ shRNA (h) Lentiviral Particles: sc-78774-V and PP2C ζ shRNA (m) Lentiviral Particles: sc-155945-V.

Molecular Weight of PP2C ζ isoforms: 55/33 kDa.


Positive Controls: RAW 264.7 whole cell lysate: sc-2211, A-431 whole cell lysate: sc-2201 or human kidney extract: sc-363764.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



PP2C ζ (C-4): sc-390214. Western blot analysis of PP2C ζ expression in NAMALWA (A), BJAB (B), Raji (C) and RAW 264.7 (D) whole cell lysates.



PP2C ζ (C-4): sc-390214. Western blot analysis of PP2C ζ expression in A-431 whole cell lysate (A) and human kidney tissue extract (B).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.