

Synaptotagmin VI (A-12): sc-390321

BACKGROUND

Synaptotagmins are a large gene family of synaptic vesicle type III integral membrane proteins that function as regulators of both exocytosis and endocytosis and are involved in neurotransmitter secretion from small secretory vesicles. Calcium binds to Synaptotagmin I which triggers neurotransmitter release at the synapse. Synaptotagmin II is phosphorylated by WNK1 in a process that regulates calcium-dependent interactions. Synaptotagmin III is involved in calcium-dependent exocytosis of secretory vesicles in endocrine cells and neurons. Synaptotagmin IV is expressed in neuronal tissues, and has the highest mRNA levels in the hippocampus. The proximity of the Synaptotagmin IV gene to markers of several psychiatric disorders suggest an involvement of synaptotagmin IV in human disease. Synaptotagmin V is a dense-core vesicle-specific protein that regulates a specific type of calcium-regulated secretion. Synaptotagmin VI interacts with adaptor protein-2 in a calcium-independent manner. Synaptotagmin VII is widely expressed in non-neuronal tissues.

REFERENCES

- Hilbush, B.S. and Morgan, J.I. 1994. A third Synaptotagmin gene, Syt3, in the mouse. Proc. Natl. Acad. Sci. USA 91: 8195-8199.
- Li, C., et al. 1995. Ca²⁺-dependent and -independent activities of neural and non-neural synaptotagmins. Nature 375: 594-599.
- Kishore, B.K., et al. 1998. Expression of Synaptotagmin VIII in rat kidney. Am. J. Physiol. 275: F131-F142.
- Xi, D., et al. 1999. Analysis of Synaptotagmin I-IV messenger RNA expression and developmental regulation in the rat hypothalamus and pituitary. Neuroscience 88: 425-435.

CHROMOSOMAL LOCATION

Genetic locus: SYT6 (human) mapping to 1p13.2; Syt6 (mouse) mapping to 3 F2.2.

SOURCE

Synaptotagmin VI (A-12) is a mouse monoclonal antibody raised against amino acids 84-137 mapping near the N-terminus of Synaptotagmin VI of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Synaptotagmin VI (A-12) is available conjugated to agarose (sc-390321 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390321 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390321 PE), fluorescein (sc-390321 FITC), Alexa Fluor® 488 (sc-390321 AF488), Alexa Fluor® 546 (sc-390321 AF546), Alexa Fluor® 594 (sc-390321 AF594) or Alexa Fluor® 647 (sc-390321 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-390321 AF680) or Alexa Fluor® 790 (sc-390321 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Synaptotagmin VI (A-12) is recommended for detection of Synaptotagmin VI of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Synaptotagmin VI siRNA (h): sc-76618, Synaptotagmin VI siRNA (m): sc-76619, Synaptotagmin VI shRNA Plasmid (h): sc-76618-SH, Synaptotagmin VI shRNA Plasmid (m): sc-76619-SH, Synaptotagmin VI shRNA (h) Lentiviral Particles: sc-76618-V and Synaptotagmin VI shRNA (m) Lentiviral Particles: sc-76619-V.

Molecular Weight (observed) of Synaptotagmin VI: 74 kDa.

Molecular Weight (predicted) of Synaptotagmin VI isoforms 1/2: 57/48 kDa.


Positive Controls: Synaptotagmin VI (h): 293T Lysate: sc-115895.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.
- Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Synaptotagmin VI (A-12): sc-390321. Western blot analysis of Synaptotagmin VI expression in non-transfected: sc-117752 (**A**) and human Synaptotagmin VI transfected: sc-115895 (**B**) 293T whole cell lysates.



Synaptotagmin VI (A-12): sc-390321. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic staining of hepatocytes.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.