

GlcAT-I (H-5): sc-390526

BACKGROUND

GlcAT-I (glucuronosyltransferase-I), also known as β -1,3-glucuronyltransferase 3 (B3GAT3), is a 335 amino acid single-pass type II membrane protein belonging to the glycosyltransferase 43 family. By using manganese as a cofactor, GlcAT-I catalyzes the formation of the glycosaminoglycan-protein linkage by way of a glucuronyl transfer reaction that is present in the final step of the biosynthesis of the linkage region of proteoglycans. Present as a disulfide-linked homodimer, GlcAT-I shows strict specificity for Gal- β -1,3-Gal- β -1,4-Xyl. Ubiquitously expressed, GlcAT-I is N-glycosylated and is localized to the Golgi apparatus membrane.

REFERENCES

1. Kitagawa, H., et al. 1998. Molecular cloning and expression of glucuronyltransferase I involved in the biosynthesis of the glycosaminoglycan-protein linkage region of proteoglycans. *J. Biol. Chem.* 273: 6615-6618.
2. Tone, Y., et al. 1999. Characterization of recombinant human glucuronyltransferase I involved in the biosynthesis of the glycosaminoglycan-protein linkage region of proteoglycans. *FEBS Lett.* 459: 415-420.
3. Ouzzine, M., et al. 2000. Structure/function of the human Gal1 β 1,3-glucuronosyltransferase. Dimerization and functional activity are mediated by two crucial cysteine residues. *J. Biol. Chem.* 275: 28254-28260.
4. Pedersen, L.C., et al. 2000. Heparan/chondroitin sulfate biosynthesis. Structure and mechanism of human glucuronyltransferase I. *J. Biol. Chem.* 275: 34580-34585.
5. Gulberti, S., et al. 2003. The functional glycosyltransferase signature sequence of the human β 1,3-glucuronosyltransferase is a XDD motif. *J. Biol. Chem.* 278: 32219-32226.
6. Venkatesan, N., et al. 2004. Stimulation of proteoglycan synthesis by glucuronosyltransferase-I gene delivery: a strategy to promote cartilage repair. *Proc. Natl. Acad. Sci. USA* 101: 18087-18092.

CHROMOSOMAL LOCATION

Genetic locus: B3GAT3 (human) mapping to 11q12.3; B3gat3 (mouse) mapping to 19 A.

SOURCE

GlcAT-I (H-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 319-334 of GlcAT-I of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-390526 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GlcAT-I (H-5) is recommended for detection of GlcAT-I of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GlcAT-I siRNA (h): sc-96989, GlcAT-I siRNA (m): sc-145416, GlcAT-I shRNA Plasmid (h): sc-96989-SH, GlcAT-I shRNA Plasmid (m): sc-145416-SH, GlcAT-I shRNA (h) Lentiviral Particles: sc-96989-V and GlcAT-I shRNA (m) Lentiviral Particles: sc-145416-V.


Molecular Weight of GlcAT-I: 37 kDa.

Positive Controls: GlcAT-I (h): 293T Lysate: sc-370997, CCRF-CEM cell lysate: sc-2225 or SK-N-SH cell lysate: sc-2410.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



GlcAT-I (H-5): sc-390526. Western blot analysis of GlcAT-I expression in CCRF-CEM (A), MOLT-4 (B) and SUP-T1 (C) whole cell lysates.



GlcAT-I (H-5): sc-390526. Western blot analysis of GlcAT-I expression in non-transfected 293T: sc-117752 (A), human GlcAT-I transfected 293T: sc-370997 (B) and SK-N-SH (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.