

XylT-I (E-5): sc-390671

BACKGROUND

Xylosyltransferase-I (XylT-I), also designated UDP-D-xylose:proteoglycan core protein β -D-xylosyltransferase 1, is a glycoprotein that catalyzes the transfer of UDP-xylose to serine residues within XT recognition sequences of target proteins. Addition of xylose to the core protein is a requirement for the biosynthesis of the glycosaminoglycan chains that are characteristic of proteoglycans. Xylosyltransferase proteins, which can be secreted, display activity in sternal cartilage chondrocytes, chondrosarcoma, nasal septum tumor and choriocarcinoma cells. XylT-I is widely expressed, with higher levels of expression detected in placenta, kidney and pancreas, and lower levels of expression observed in skeletal muscle. Xylosyltransferase-II (XylT-II), also designated UDP-D-xylose:proteoglycan core protein β -D-xylosyltransferase 2, is also widely expressed, with higher levels of expression detected in kidney and pancreas.

REFERENCES

1. Götting, C., et al. 2004. Analysis of the DXD motifs in human xylosyltransferase I required for enzyme activity. *J. Biol. Chem.* 279: 42566-42573.
2. Schön, S., et al. 2005. Impact of polymorphisms in the genes encoding xylosyltransferase I and a homologue in type 1 diabetic patients with and without nephropathy. *Kidney Int.* 68: 1483-1490.
3. Götting, C., et al. 2005. Elevated xylosyltransferase I activities in pseudoxanthoma elasticum (PXE) patients as a marker of stimulated proteoglycan biosynthesis. *J. Mol. Med.* 83: 984-992.
4. Kuhn, J., et al. 2005. Xylosyltransferase I acceptor properties of fibroblast growth factor and its fragment bFGF (1-24). *Biochem. Biophys. Res. Commun.* 333: 156-166.
5. Schöttler, M., et al. 2005. Serum not affected by renal insufficiency. *Clin. Biochem.* 38: 486-488.

CHROMOSOMAL LOCATION

Genetic locus: XYLT1 (human) mapping to 16p12.3; Xylt1 (mouse) mapping to 7 F2.

SOURCE

XylT-I (E-5) is a mouse monoclonal antibody raised against amino acids 871-944 mapping near the C-terminus of XylT-I of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

XylT-I (E-5) is available conjugated to agarose (sc-390671 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390671 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390671 PE), fluorescein (sc-390671 FITC), Alexa Fluor® 488 (sc-390671 AF488), Alexa Fluor® 546 (sc-390671 AF546), Alexa Fluor® 594 (sc-390671 AF594) or Alexa Fluor® 647 (sc-390671 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-390671 AF680) or Alexa Fluor® 790 (sc-390671 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

XylT-I (E-5) is recommended for detection of XylT-I of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for XylT-I siRNA (h): sc-61817, XylT-I siRNA (m): sc-61818, XylT-I shRNA Plasmid (h): sc-61817-SH, XylT-I shRNA Plasmid (m): sc-61818-SH, XylT-I shRNA (h) Lentiviral Particles: sc-61817-V and XylT-I shRNA (m) Lentiviral Particles: sc-61818-V.

Molecular Weight of XylT-I: 72 kDa.


Positive Controls: HEL 92.1.7 cell lysate: sc-2270, JAR cell lysate: sc-2276 or MIA PaCa-2 cell lysate: sc-2285.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



XylT-I (E-5): sc-390671. Western blot analysis of XylT-I expression in JAR (A), HEL 92.1.7 (B), Ramos (C) and HEK293 (D) whole cell lysates.



XylT-I (E-5): sc-390671. Western blot analysis of XylT-I expression in JAR (A) and MIA PaCa-2 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA