

Synapsin Ia/b (D-8): sc-390867

BACKGROUND

Synapsins are synaptic vesicle-associated phosphoproteins that regulate synaptic vesicle exocytosis and may be involved in synaptogenesis. Evidence suggests that Synapsin I, Synapsin II and Synapsin IIIa are ATP-binding proteins that are regulated by Ca^{2+} and calmodulin binding. Ca^{2+} has been shown to stimulate ATP binding to Synapsin I, to have no effect on Synapsin II and to inhibit Synapsin III. Synapsin I and Synapsin II both undergo alternative splicing to produce two forms of each protein, Synapsin Ia and Ib and Synapsin IIa and IIb, respectively. Synapsin III gives rise to at least three isoforms: Synapsin IIIa, IIIb and IIIc. Synapsin III plays unique roles both in early axon outgrowth and in the regulation of synaptic vesicle trafficking. In cultured mouse hippocampal neurons, Synapsin III is expressed early during development, with levels peaking seven days after plating and declining thereafter. Synapsin III is highly concentrated in growth cones.

REFERENCES

1. Sudhof, T.C., et al. 1989. Synapsins: mosaics of shared and individual domains in a family of synaptic vesicle phosphoproteins. *Science* 245: 1474-1480.
2. Sudhof, T.C. 1990. The structure of the human synapsin I gene and protein. *J. Biol. Chem.* 265: 7849-7852.
3. Melloni, R.H., Jr., et al. 1994. Temporal onset of synapsin I gene expression coincides with neuronal differentiation during the development of the nervous system. *J. Comp. Neurol.* 342: 449-462.
4. Nicol, S., et al. 1997. Ca^{2+} -dependent interaction with calmodulin is conserved in the synapsin family: identification of a high-affinity site. *Biochemistry* 36: 11487-11495.
5. Hosaka, M. and Sudhof, T.C. 1998. Synapsins I and II are ATP-binding proteins with differential Ca^{2+} regulation. *J. Biol. Chem.* 273: 1425-1429.

CHROMOSOMAL LOCATION

Genetic locus: SYN1 (human) mapping to Xp11.23; Syn1 (mouse) mapping to X A1.3.

SOURCE

Synapsin Ia/b (D-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 651-665 near the C-terminus of Synapsin Ia of human origin.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-390867 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Synapsin Ia/b (D-8) is recommended for detection of Synapsin Ia and Synapsin Ib of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Synapsin Ia/b siRNA (h): sc-37012, Synapsin Ia/b siRNA (m): sc-37013, Synapsin Ia/b shRNA Plasmid (h): sc-37012-SH, Synapsin Ia/b shRNA Plasmid (m): sc-37013-SH, Synapsin Ia/b shRNA (h) Lentiviral Particles: sc-37012-V and Synapsin Ia/b shRNA (m) Lentiviral Particles: sc-37013-V.

Molecular Weight of Synapsin Ia: 80 kDa.


Molecular Weight of Synapsin Ib: 86 kDa.

Positive Controls: mouse brain extract: sc-2253 or rat cerebellum extract: sc-2398.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



Synapsin Ia/b (D-8): sc-390867. Western blot analysis of Synapsin Ia/b expression in mouse brain (A) and rat cerebellum (B) tissue extracts.

SELECT PRODUCT CITATIONS

1. Gassowska-Dobrowolska, M., et al. 2020. Prenatal exposure to valproic acid affects microglia and synaptic ultrastructure in a brain-region-specific manner in young-adult male rats: relevance to autism spectrum disorders. *Int. J. Mol. Sci.* 21: 3576.

RESEARCH USE

For research use only, not for use in diagnostic procedures.