

# AChR $\delta$ (C-4): sc-390896

## BACKGROUND

Members of the ligand-gated ion channel receptor family are characterized by their fast transmitting response to neurotransmitters. Two important members of this family are the nicotinic acetylcholine and glutamate receptors, both of which are composed of five homologous subunits forming a transmembrane aqueous pore. These transmembrane receptors change conformation in response to their cognate neurotransmitter. Nicotinic acetylcholine receptors (AChRs) are found at the postsynaptic membrane of the neuromuscular junction and bind acetylcholine molecules, allowing ions to move through the pore. Glutamate receptors are found in the postsynaptic membrane of cells in the central nervous system. The activity that is generated at the synapse by the binding of acetylcholine is terminated by acetylcholinesterase, an enzyme that rapidly hydrolyzes acetylcholine. AChR $\delta$ , also known as CMS2A, FCCMS, SCCMS or CHRND, is a 517 amino acid multi-pass membrane protein that is associated with lethal type multiple pterygium syndrome, congenital myasthenic syndrome slow-channel type (SCCMS) and congenital myasthenic syndrome fast-channel type (FCCMS).

## CHROMOSOMAL LOCATION

Genetic locus: CHRND (human) mapping to 2q37.1; Chrnd (mouse) mapping to 1 D.

## SOURCE

AChR $\delta$  (C-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 349-502 near the C-terminus of AChR $\delta$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

AChR $\delta$  (C-4) is available conjugated to agarose (sc-390896 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390896 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390896 PE), fluorescein (sc-390896 FITC), Alexa Fluor<sup>®</sup> 488 (sc-390896 AF488), Alexa Fluor<sup>®</sup> 546 (sc-390896 AF546), Alexa Fluor<sup>®</sup> 594 (sc-390896 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-390896 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-390896 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-390896 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-390896 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

AChR $\delta$  (C-4) is recommended for detection of AChR $\delta$  of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for AChR $\delta$  siRNA (h): sc-42540, AChR $\delta$  siRNA (m): sc-42541, AChR $\delta$  shRNA Plasmid (h): sc-42540-SH, AChR $\delta$  shRNA Plasmid (m): sc-42541-SH, AChR $\delta$  shRNA (h) Lentiviral Particles: sc-42540-V and AChR $\delta$  shRNA (m) Lentiviral Particles: sc-42541-V.


Molecular Weight of AChR $\delta$ : 60 kDa.

Positive Controls: SJRH30 cell lysate: sc-2287 or BC<sub>3</sub>H1 cell lysate: sc-2299.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG $\kappa$  BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



AChR $\delta$  (C-4): sc-390896. Western blot analysis of AChR $\delta$  expression in SJRH30 (A) and BC<sub>3</sub>H1 (B) whole cell lysates.



AChR $\delta$  (C-4): sc-390896. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing membrane staining of myocytes.

## SELECT PRODUCT CITATIONS

1. Condorelli, R.A., et al. 2017. Nicotine effects and receptor expression on human spermatozoa: possible neuroendocrine mechanism. *Front. Physiol.* 8: 177.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.