



PP2B-A γ siRNA (m): sc-39198

BACKGROUND

In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases. In general, the protein phosphatase (PP) holoenzyme is a trimeric complex composed of a regulatory subunit, a variable subunit and a catalytic subunit. Four major families of protein phosphatase catalytic subunit have been identified, designated PP1, PP2A, PP2B and PP2C. An additional protein phosphatase catalytic subunit, PPX (also known as PP4), is a putative member of a novel PP family. The PP2B family comprises subfamily members PP2B-A α , PP2B-A β and PP2B-A γ . Two additional regulatory subunits been identified, designated PP2B-B1 and PP2B-B2.

REFERENCES

1. Ueki, K., et al. 1992. Structure and expression of two isoforms of the murine calmodulin-dependent protein phosphatase regulatory subunit (calcineurin B). *Biochem. Biophys. Res. Commun.* 187: 537-543.
2. Mumby, M.C., et al. 1993. Protein serine/threonine phosphatases: structure, regulation, and functions in cell growth. *Phys. Rev.* 73: 673-699.
3. Hendrix, P., et al. 1993. Structure and expression of a 72-kDa regulatory subunit of protein phosphatase 2A. Evidence for different size forms produced by alternative splicing. *J. Biol. Chem.* 268: 15267-15276.
4. Cohen, P.T. 1993. Important roles for novel protein phosphatases dephosphorylating serine and threonine residues. *Biochem. Soc. Trans.* 21: 884-888.
5. Okubo, S., et al. 1994. A regulatory subunit of smooth muscle Myosin bound phosphatase. *Biochem. Biophys. Res. Commun.* 200: 429-434.
6. Wera, S., et al. 1995. Serine/threonine protein phosphatases. *Biochem. J.* 311: 17-29.
7. Lohse, D.L., et al. 1995. Insights derived from the structures of the Ser/Thr phosphatases calcineurin and protein phosphatase 1. *Structure* 3: 987-990.

CHROMOSOMAL LOCATION

Genetic locus: Ppp3cc (mouse) mapping to 14 D2.

PRODUCT

PP2B-A γ siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PP2B-A γ shRNA Plasmid (m): sc-39198-SH and PP2B-A γ shRNA (m) Lentiviral Particles: sc-39198-V as alternate gene silencing products.

For independent verification of PP2B-A γ (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-39198A, sc-39198B and sc-39198C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PP2B-A γ siRNA (m) is recommended for the inhibition of PP2B-A γ expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PP2B-A γ gene expression knockdown using RT-PCR Primer: PP2B-A γ (m)-PR: sc-39198-PR (20 μ l, 418 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.