

Mos^{x_e} (xQ-20): sc-392290

BACKGROUND

The cellular homolog of the v-Mos oncogene, originally derived from the Moloney strain murine sarcoma virus, has now been isolated from several species including human, mouse, chicken and *Xenopus*. In each of these species, the c-Mos proto-oncogene locus consists of a single exon which encodes Mos, a serine/threonine protein kinase involved in the regulation of the meiotic cell cycle in vertebrate oocytes. Mos function during oocyte maturation is thought to be exerted through the stabilization or activation of the maturation promoting factor (MPF) complex. While the level of Mos expression is generally low in somatic cells, Mos is expressed at relatively high levels in germ cells. Mos can efficiently cause oncogenic transformation of somatic cells when ectopically expressed during the G₁ phase of the cell cycle. Mos activates MAPK, possibly through direct phosphorylation of the MAPKK MEK-1.

REFERENCES

1. Klein, G., et al. 1981. The role of gene dosage and genetic transpositions in carcinogenesis. *Nature* 294: 313-318.
2. Watson, R., et al. 1982. Human DNA sequence homologous to the transforming gen (Mos) of Moloney murine sarcoma virus. *Proc. Natl. Acad. Sci. USA* 79: 4078-4082.
3. Propst, F., et al. 1985. Expression of c-Mos proto-oncogene transcripts in mouse tissues. *Nature* 315: 516-518.
4. Schmidt, M., et al. 1988. Chicken homolog of the Mos proto-oncogene. *Mol. Cell. Biol.* 8: 923-929.
5. Okazaki, K., et al. 1995. MAP kinase activation is essential for oncogenic transformation of NIH/3T3 cells by Mos. *Oncogene* 10: 1149-1157.
6. Chen, M., et al. 1995. Ser-3 is important for regulating Mos interaction with and stimulation of mitogen-activated protein kinase kinase. *Mol. Cell. Biol.* 15: 4727-4734.
7. Pham, C.D., et al. 1995. Characterization of MEK-1 phosphorylation by the v-Mos protein. *Oncogene* 10: 1683-1688.
8. Robertson, S.C., et al. 1996. Identification of an autoinhibitory region in the activation loop of the Mos protein kinase. *Mol. Cell. Biol.* 16: 3472-3479.
9. Mendez, R., et al. 2000. Phosphorylation of CPE binding factor by Eg2 regulates translation of c-Mos mRNA. *Nature* 404: 302-307.

SOURCE

Mos^{x_e} (xQ-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Mos^{x_e} of *Xenopus laevis* origin.

PRODUCT

Each vial contains 200 µg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Mos^{x_e} (B-4) is recommended for detection of Mos p39 of *Xenopus laevis* origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of Mos^{x_e}: 37 kDa.

Positive Controls: XLK-WG whole cell lysate: sc-364801.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Mos^{x_e} (xQ-20): sc-392290. Western blot analysis of Mos^{x_e} expression in XLK-WG whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.