

Lutropin β siRNA (m): sc-39320

BACKGROUND

Various hormones are secreted from the anterior pituitary gland during development and growth. Lutropin, also called luteinizing hormone (LH), plays a role in spermatogenesis and ovulation by stimulating the testes and ovaries to produce steroids. LH, like many of the anterior pituitary hormones, consists of heterodimers formed from a common α chain and a unique β chain. Lutropin exists in a variety of isoforms, as the hormone is proteolytically processed and metabolized throughout circulation. LH modulates the processing of Amyloid- β precursor protein and Amyloid- β deposition. Pituitary exit of LH and FSH occur via different secretion pathways, and are released spatially from the pituitary via different circulatory routes.

REFERENCES

1. Couzinet, B., et al. 1993. The control of gonadotrophin secretion by ovarian steroids. *Hum. Reprod.* 2: 97-101.
2. Birken, S., et al. 1996. Metabolism of hCG and hLH to multiply urinary forms. *Mol. Cell. Endocrinol.* 125: 121-131.
3. Sherman, G.B., et al. 1997. Characterization and phylogenetic significance of rhinoceros luteinizing hormone β (LH β) subunit messenger RNA structure, complementary DNA sequence and gene copy number. *Gene* 195: 131-139.
4. Hakola, K., et al. 1998. Recombinant forms of rat and human luteinizing hormone and follicle-stimulating hormone; comparison of functions *in vitro* and *in vivo*. *J. Endocrinol.* 158: 441-448.
5. Arnold, C.J., et al. 1998. The human follitropin α subunit C-terminus collaborates with a β subunit cystine noose and an α subunit loop to assemble a receptor-binding domain competent for signal transduction. *Biochemistry* 37: 1762-1768.
6. Jablonka-Shariff, A., et al. 2002. Evolution of Lutropin to chorionic gonadotropin generates a specific routing signal for apical release *in vivo*. *J. Biol. Chem.* 277: 879-882.
7. Xing, Y., et al. 2004. Use of protein knobs to characterize the position of conserved α subunit regions in Lutropin receptor complexes. *J. Biol. Chem.* 279: 44427-44437.
8. Bernard, M.P., et al. 2004. Only a portion of the small seatbelt loop in human choriogonadotropin appears capable of contacting the Lutropin receptor. *J. Biol. Chem.* 279: 44438-44441.

CHROMOSOMAL LOCATION

Genetic locus: Lhb (mouse) mapping to 7 B4.

PRODUCT

Lutropin β siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Lutropin β shRNA Plasmid (m): sc-39320-SH and Lutropin β shRNA (m) Lentiviral Particles: sc-39320-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Lutropin β siRNA (m) is recommended for the inhibition of Lutropin β expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Lutropin β (C-6): sc-373941 is recommended as a control antibody for monitoring of Lutropin β gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Lutropin β gene expression knockdown using RT-PCR Primer: Lutropin β (m)-PR: sc-39320-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.