

copine 1/3 (E-8): sc-393249

BACKGROUND

The copine family is composed of evolutionarily conserved, calcium-dependent membrane-binding proteins. Members of the copine family are involved in signal transduction and membrane trafficking, and are capable of binding phospholipids in a calcium-dependent manner. Copine 1, also known as CPNE1, CPN1 or COPN1, is ubiquitously expressed and contains two N-terminal C2 domains and one C-terminal VWFA (von Willebrand factor A) domain, which is also referred to as the A domain or the core domain. Via its VWFA domain, copine 1 directly interacts with the ubiquitin-conjugating enzyme UBC12 and may play a role in the regulation of TNF α -signaling. Copine 2, also known as COPN2 or CPNE2, contains two C2 domains and one C-terminal VWFA domain. Copine 3, also known as CPNE3, CPN3 or COPN3, is ubiquitously expressed and contains two N-terminal C2 domains and one C-terminal VWFA domain. Differing from other members of the copine family, copine 3 may possess some intrinsic kinase activity.

REFERENCES

- Creutz, C.E., et al. 1998. The copines, a novel class of C2 domain-containing, calcium-dependent, phospholipid-binding proteins conserved from *Paramecium* to humans. *J. Biol. Chem.* 273: 1393-1402.
- Tomsig, J.L. and Creutz, C.E. 2000. Biochemical characterization of copine: a ubiquitous Ca²⁺-dependent, phospholipid-binding protein. *Biochemistry* 39: 16163-16175.
- Jambunathan, N., et al. 2001. A humidity-sensitive *Arabidopsis* copine mutant exhibits precocious cell death and increased disease resistance. *Plant Cell* 13: 2225-2240.
- Cowland, J.B., et al. 2003. Tissue expression of copines and isolation of copines I and III from the cytosol of human neutrophils. *J. Leukoc. Biol.* 74: 379-388.

CHROMOSOMAL LOCATION

Genetic locus: CPNE1 (human) mapping to 20q11.22, CPNE3 (human) mapping to 8q21.3; Cpn1 (mouse) mapping to 2 H1, Cpn3 (mouse) mapping to 4 A3.

SOURCE

copine 1/3 (E-8) is a mouse monoclonal antibody raised against amino acids 266-346 mapping within an internal region of copine I of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

copine 1/3 (E-8) is available conjugated to agarose (sc-393249 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-393249 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-393249 PE), fluorescein (sc-393249 FITC), Alexa Fluor[®] 488 (sc-393249 AF488), Alexa Fluor[®] 546 (sc-393249 AF546), Alexa Fluor[®] 594 (sc-393249 AF594) or Alexa Fluor[®] 647 (sc-393249 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-393249 AF680) or Alexa Fluor[®] 790 (sc-393249 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

copine 1/3 (E-8) is recommended for detection of copine 1 and copine 3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with copine 2.


Molecular Weight of copine 1/3: 60 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, CCRF-CEM cell lysate: sc-2225 or SK-BR-3 cell lysate: sc-2218.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



copine 1/3 (E-8): sc-393249. Western blot analysis of copine 1/3 expression in Hep G2 (A), SK-BR-3 (B), U-937 (C) and CCRF-CEM (D) whole cell lysates.



copine 1/3 (E-8): sc-393249. Immunofluorescence staining of formalin-fixed Hep G2 (A) and SW480 (B) cells showing cytoplasmic and nuclear localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.