

HYLS1 (F-12): sc-393492

BACKGROUND

The hydrolethalus syndrome protein 1 (HYLS1) is a widely conserved protein that plays an essential role in cilia formation. A single amino acid mutation in the HYLS1 gene leads to a perinatal lethal disorder termed hydrolethalus syndrome, a severe fetal malformation syndrome characterized by central nervous system (CNS) malformation such as hydrocephaly and absent midline structures of the brain, micrognathia, defective lobation of the lungs and polydactyly. The gene encoding HYLS1 maps to human chromosome 11, which makes up around 4% of human genomic DNA and is considered a gene and disease association dense chromosome. The chromosome 11 encoded Atm gene is important for regulation of cell cycle arrest and apoptosis following double strand DNA breaks. Atm mutation leads to the disorder known as ataxia-telangiectasia.

REFERENCES

- Grossfeld, P.D., et al. 2004. The 11q terminal deletion disorder: a prospective study of 110 cases. *Am. J. Med. Genet. A* 129A: 51-61.
- Mee, L., et al. 2005. Hydrolethalus syndrome is caused by a missense mutation in a novel gene HYLS1. *Hum. Mol. Genet.* 14: 1475-1488.
- Taylor, T.D., et al. 2006. Human chromosome 11 DNA sequence and analysis including novel gene identification. *Nature* 440: 497-500.
- Lee, S.P., et al. 2007. Phase I study of eptifibatid in patients with sickle cell anaemia. *Br. J. Haematol.* 139: 612-620.
- Lee, J.H., et al. 2007. Activation and regulation of ATM kinase activity in response to DNA double-strand breaks. *Oncogene* 26: 7741-7748.
- Paetau, A., et al. 2008. Hydrolethalus syndrome: neuropathology of 21 cases confirmed by HYLS1 gene mutation analysis. *J. Neuropathol. Exp. Neurol.* 67: 750-762.
- Kaste, S.C., et al. 2008. Wilms tumour: prognostic factors, staging, therapy and late effects. *Pediatr. Radiol.* 38: 2-17.
- Dammermann, A., et al. 2009. The hydrolethalus syndrome protein HYLS-1 links core centriole structure to cilia formation. *Genes Dev.* 23: 2046-2059.

CHROMOSOMAL LOCATION

Genetic locus: HYLS1 (human) mapping to 11q24.2; Hyls1 (mouse) mapping to 9 A4.

SOURCE

HYLS1 (F-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 38-57 within an internal region of HYLS1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₃ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-393492 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

HYLS1 (F-12) is recommended for detection of HYLS1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HYLS1 siRNA (h): sc-96710, HYLS1 shRNA Plasmid (h): sc-96710-SH and HYLS1 shRNA (h) Lentiviral Particles: sc-96710-V.


Molecular Weight of HYLS1: 40 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409 or SK-MEL-24 whole cell lysate: sc-364259.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



HYLS1 (F-12): sc-393492. Western blot analysis of HYLS1 expression in IMR-32 (A) and SK-MEL-24 (B) whole cell lysates.



HYLS1 (F-12): sc-393492. Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing nuclear staining of cells in semi-iferous ducts and cytoplasmic staining of Leydig cells.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.