

Thrombospondin 1 (C-9): sc-393503

BACKGROUND

The thrombospondin proteins (TSP 1-4) compose a family of glycoproteins that are involved in cell-to-cell and cell-to-matrix signaling. These extracellular, cell-surface proteins form complexes of both homo- and heteromultimers. Thrombospondins play a role in development, aggregation of platelets, adhesion and migration of cells, and progression of cells through the growth cycle. Thrombospondin 1 is released from platelets in response to Thrombin stimulation and is a transient component of the extracellular matrix of developing and repairing tissues. Thrombospondin 2 shares a high degree of homology with Thrombospondin 1 and is thought to have overlapping but unique functions. Thrombospondin 3 is a developmentally regulated heparin binding protein. Thrombospondin 4 is neuronally expressed and stimulates neurite outgrowth.

REFERENCES

1. Mosher, D.F. 1990. Physiology of thrombospondin. *Annu. Rev. Med.* 41: 85-97.
2. Bornstein, P., et al. 1991. A second, expressed thrombospondin gene (Thbs2) exists in the mouse genome. *J. Biol. Chem.* 266: 12821-12824.
3. LaBell, T.L., et al. 1992. Thrombospondin II: partial cDNA sequence, chromosome location and expression of a second member of the thrombospondin gene family in humans. *Genomics* 12: 421-429.
4. O'Rourke, K.M., et al. 1992. Thrombospondin 1 and Thrombospondin 2 are expressed as both homo- and heterotrimers. *J. Biol. Chem.* 267: 24921-24924.
5. Jahav, J. 1993. The functions of thrombospondin and its involvement in physiology and pathophysiology. *Biochem. Biophys. Acta* 1182: 1-14.

CHROMOSOMAL LOCATION

Genetic locus: THBS1 (human) mapping to 15q14; Thbs1 (mouse) mapping to 2 E5.

SOURCE

Thrombospondin 1 (C-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 25-48 near the N-terminus of Thrombospondin 1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-393503 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Thrombospondin 1 (C-9) is recommended for detection of Thrombospondin 1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Thrombospondin 1 (C-9) is also recommended for detection of Thrombospondin 1 in additional species, including equine, bovine and porcine.


Suitable for use as control antibody for Thrombospondin 1 siRNA (h): sc-36665, Thrombospondin 1 siRNA (m): sc-36666, Thrombospondin 1 siRNA (r): sc-270413, Thrombospondin 1 shRNA Plasmid (h): sc-36665-SH, Thrombospondin 1 shRNA Plasmid (m): sc-36666-SH, Thrombospondin 1 shRNA Plasmid (r): sc-270413-SH, Thrombospondin 1 shRNA (h) Lentiviral Particles: sc-36665-V, Thrombospondin 1 shRNA (m) Lentiviral Particles: sc-36666-V and Thrombospondin 1 shRNA (r) Lentiviral Particles: sc-270413-V.

Molecular Weight of Thrombospondin 1 various forms: 165-198 kDa.


Molecular Weight of Thrombospondin 1 homotrimer: 420 kDa.

Positive Controls: BJ whole cell lysate: sc-364359, CCD-1064Sk cell lysate: sc-2263 or HUV-EC-C whole cell lysate: sc-364180.

DATA



Thrombospondin 1 (C-9): sc-393503. Western blot analysis of Thrombospondin 1 expression in CCD-1064Sk (A), BJ (B), HUV-EC-C (C) and MDA-MB-231 (D) whole cell lysates and human pancreas tissue extract (E).



Thrombospondin 1 (C-9): sc-393503. Western blot analysis of Thrombospondin 1 expression in NCI-H292 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Anastasi, C., et al. 2020. BMP-1 disrupts cell adhesion and enhances TGF-β activation through cleavage of the matricellular protein Thrombospondin-1. *Sci. Signal.* 13: eaba3880.
2. Hellewell, A.L., et al. 2022. PDIA3/ERp57 promotes a matrix-rich secretome that stimulates fibroblast adhesion through CCN2. *Am. J. Physiol. Cell. Physiol.* 322: C624-C644.



See **Thrombospondin 1 (A6.1): sc-59887** for additional antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.