

# HSF1 (C-9): sc-393509

## BACKGROUND

Prokaryotic and eukaryotic cells respond to thermal and chemical stress by inducing a group of genes collectively designated heat shock genes. In eukaryotes, this gene expression is regulated primarily at the transcription level. Heat shock transcription factors 1 and 2 (HSF1 and HSF2), also designated HSTF1 and HSTF2, are involved in this regulation. HSF1 and HSF2 are upregulated by estrogen at both the mRNA and protein level. HSF1 is normally found as a monomer, whose transcriptional activity is repressed by constitutive phosphorylation. Upon activation, HSF1 forms trimers, gains DNA binding activity and is translocated to the nucleus. HSF2 activity is associated with differentiation and development and, like HSF1, binds DNA as a trimer. Both HSF1 and HSF2 are known to be induced by proteasome inhibitors of the ubiquitin pathway.

## CHROMOSOMAL LOCATION

Genetic locus: HSF1 (human) mapping to 8q24.3; Hsf1 (mouse) mapping to 15 D3.

## SOURCE

HSF1 (C-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 508-529 at the C-terminus of HSF1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-393509 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-393509 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## APPLICATIONS

HSF1 (C-9) is recommended for detection of HSF1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HSF1 (C-9) is also recommended for detection of HSF1 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for HSF1 siRNA (h): sc-35611, HSF1 siRNA (m): sc-35612, HSF1 siRNA (r): sc-270440, HSF1 shRNA Plasmid (h): sc-35611-SH, HSF1 shRNA Plasmid (m): sc-35612-SH, HSF1 shRNA Plasmid (r): sc-270440-SH, HSF1 shRNA (h) Lentiviral Particles: sc-35611-V, HSF1 shRNA (m) Lentiviral Particles: sc-35612-V and HSF1 shRNA (r) Lentiviral Particles: sc-270440-V.

HSF1 (C-9) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of HSF1: 89-90 kDa.

Positive Controls: HSF1 (h): 293T Lysate: sc-171930.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



HSF1 (C-9): sc-393509. Western blot analysis of HSF1 expression in Hep G2 (A), MCF7 (B), A2058 (C), K-562 (D) and HeLa (E) whole cell lysates.



HSF1 (C-9): sc-393509. Western blot analysis of HSF1 expression in non-transfected: sc-117752 (A) and human HSF1 transfected: sc-171930 (B) 293T whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.



See **HSF1 (E-4): sc-17757** for HSF1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.