


## COL1A2 (E-6): sc-393573



The Power to Question

## BACKGROUND

The extensive family of COL gene products (collagens) is composed of several chain types, including fibril-forming interstitial collagens (types I, II, III and V) and basement membrane collagens (type IV), each type containing multiple isoforms. Collagens are fibrous, extracellular matrix proteins with high tensile strength and are the major components of connective tissue, such as tendons and cartilage. All collagens contain a triple helix domain and frequently show lateral self-association in order to form complex connective tissues. Several collagens also play a role in cell adhesion, important for maintaining normal tissue architecture and function.

## CHROMOSOMAL LOCATION

Genetic locus: COL1A2 (human) mapping to 7q21.3; Col1a2 (mouse) mapping to 6 A1.

## SOURCE

COL1A2 (E-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 1064-1093 at the C-terminus of Collagen  $\alpha 2$  Type I of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> lambda light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

COL1A2 (E-6) is available conjugated to agarose (sc-393573 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-393573 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-393573 PE), fluorescein (sc-393573 FITC), Alexa Fluor<sup>®</sup> 488 (sc-393573 AF488), Alexa Fluor<sup>®</sup> 546 (sc-393573 AF546), Alexa Fluor<sup>®</sup> 594 (sc-393573 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-393573 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-393573 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-393573 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-393573 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## APPLICATIONS

COL1A2 (E-6) is recommended for detection of Collagen  $\alpha 2$  Type I of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for COL1A2 siRNA (h): sc-72156, COL1A2 siRNA (m): sc-43061, COL1A2 shRNA Plasmid (h): sc-72156-SH, COL1A2 shRNA Plasmid (m): sc-43061-SH, COL1A2 shRNA (h) Lentiviral Particles: sc-72156-V and COL1A2 shRNA (m) Lentiviral Particles: sc-43061-V.


Molecular Weight of COL1A2 precursor: 130-140 kDa.

Molecular Weight of mature COL1A2: 70-90 kDa.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



COL1A2 (E-6) HRP: sc-393573 HRP. Direct western blot analysis of COL1A2 expression in 3T3-L1 (A), CCD-1064Sk (B), A-10 (C), C2C12 (D) and C3H/10T1/2 (E) whole cell lysates.



COL1A2 (E-6): sc-393573. Immunoperoxidase staining of formalin fixed, paraffin-embedded rat placenta tissue showing membrane and cytoplasmic staining of trophoblastic cells and extracellular staining of connective tissue (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded rat testis tissue showing membrane and cytoplasmic staining of cells in seminiferous ducts and cytoplasmic staining of Leydig cells (B).

## SELECT PRODUCT CITATIONS

- Miotti, S., et al. 2017. Antibody-mediated blockade of JMJD6 interaction with collagen I exerts antifibrotic and antimetastatic activities. *FASEB J.* 31: 5356-5370.
- Li, S.S., et al. 2019. Reduction of PGRN increased fibrosis during skin wound healing in mice. *Histol. Histopathol.* 34: 765-774.
- Hohl, M., et al. 2020. Cathepsin A contributes to left ventricular remodeling by degrading extracellular superoxide dismutase in mice. *J. Biol. Chem.* 295: 12605-12617.
- Bansod, S., et al. 2021. Inhibition of discoidin domain receptors by imatinib prevented pancreatic fibrosis demonstrated in experimental chronic pancreatitis model. *Sci. Rep.* 11: 12894.
- Wieczfinska, J., et al. 2022. Relaxin affects airway remodeling genes expression through various signal pathways connected with transcription factors. *Int. J. Mol. Sci.* 23: 8413.
- Piñero-Hermida, S., et al. 2022. Consequences of telomere dysfunction in fibroblasts, club and basal cells for lung fibrosis development. *Nat. Commun.* 13: 5656.
- Soundararajan, A., et al. 2022. Multiomics analysis reveals the mechanical stress-dependent changes in trabecular meshwork cytoskeletal-extracellular matrix interactions. *Front. Cell Dev. Biol.* 10: 874828.
- Qin, L., et al. 2023. Docosahexaenoic acid administration improves diabetes-induced cardiac fibrosis through enhancing fatty acid oxidation in cardiac fibroblast. *J. Nutr. Biochem.* 113: 109244.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA