

Cdk4 (F-8): sc-393653

BACKGROUND

Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key substrates involved in each phase of cell cycle progression. Another family of proteins, Cdk inhibitors, also plays a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Several Cdk proteins have been identified, including Cdk2-Cdk8, PCTAIRE-1-3, PITSLRE and PITSLRE. Cdk4, in complex with D-type cyclins, is thought to regulate cell growth during the G₁ phase of the cell cycle. This association with a D-type cyclin upregulates Cdk4 activity, whereas binding to the Cdk inhibitor p16 downregulates Cdk4 activity. Activation of the Cdk4-cyclin complexes requires phosphorylation on a single threonyl residue of Cdk4, catalyzed by a Cdk-activating protein (CAK).

CHROMOSOMAL LOCATION

Genetic locus: CDK4 (human) mapping to 12q14.1; Cdk4 (mouse) mapping to 10 D3.

SOURCE

Cdk4 (F-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 278-303 at the C-terminus of Cdk4 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-393653 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-393653 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

Cdk4 (F-8) is recommended for detection of Cdk4 p34 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Cdk4 siRNA (h): sc-29261, Cdk4 siRNA (m): sc-29262, Cdk4 shRNA Plasmid (h): sc-29261-SH, Cdk4 shRNA Plasmid (m): sc-29262-SH, Cdk4 shRNA (h) Lentiviral Particles: sc-29261-V and Cdk4 shRNA (m) Lentiviral Particles: sc-29262-V.

Cdk4 (F-8) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Cdk4: 34 kDa.

Positive Controls: Cdk4 (m): 293T Lysate: sc-119150, SK-BR-3 cell lysate: sc-2218 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Cdk4 (F-8): sc-393653. Western blot analysis of Cdk4 expression in non-transfected 293T: sc-117752 (A), mouse Cdk4 transfected 293T: sc-119150 (B), SK-BR-3 (C), HeLa (D) and Ramos (E) whole cell lysates.



Cdk4 (F-8): sc-393653. Immunoperoxidase staining of formalin fixed, paraffin-embedded human fallopian tube tissue showing cytoplasmic staining of glandular cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing cytoplasmic and membrane staining of glandular cells (B).

SELECT PRODUCT CITATIONS

- Bencivenga, D., et al. 2014. p27 Kip1 serine 10 phosphorylation determines its metabolism and interaction with cyclin-dependent kinases. *Cell Cycle* 13: 3768-3782.
- Huang, R.S., et al. 2018. MicroRNA-381 suppresses the growth and increases cisplatin sensitivity in non-small cell lung cancer cells through inhibition of nuclear factor-κB signaling. *Biomed. Pharmacother.* 98: 538-544.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **Cdk4 (DCS-35): sc-23896** for Cdk4 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.