

# hnRNP A2/B1 (C-3): sc-393674

## BACKGROUND

Heterogeneous nuclear ribonucleoproteins (hnRNPs) constitute a set of polypeptides that contribute to mRNA transcription and pre-mRNA processing as well as mature mRNA transport to the cytoplasm and translation. They also bind heterogeneous nuclear RNA (hnRNA), which are the transcripts produced by RNA polymerase II. There are approximately 20 known hnRNP proteins, and their complexes are the major constituents of the spliceosome. The majority of hnRNP proteins are localized to the nucleus; however some shuttle between the nucleus and the cytoplasm. The A/B subfamily of hnRNPs include A1, A2/B1, A3 and A0, and in *Xenopus*, hnRNP A1, A2 and A3 are ubiquitously expressed throughout development as well as in adult tissues. hnRNP A1 and A2/B1 regulate the processing of pre-mRNA by directly antagonizing the association of various splicing factors and by influencing the splice site selection on pre-mRNA. The hnRNP A0 gene is distinct from the other A/B family members, and it encodes a low-abundance protein, which is implicated in mRNA stability.

## REFERENCES

- Good, P. J., et al. 1993. Three new members of the RNP protein family in *Xenopus*. *Nucleic Acids Res.* 21: 999-1006.
- Badolato, J., et al. 1995. Identification and characterisation of a novel human RNA-binding protein. *Gene* 166: 323-337.
- Siomi, H., et al. 1995. A nuclear localization domain in the hnRNP A1 protein. *J. Cell Biol.* 129: 551-560.
- Myer, V.E., et al. 1995. Isolation and characterization of a novel, low abundance hnRNP protein: A0. *RNA* 1: 171-182.

## CHROMOSOMAL LOCATION

Genetic locus: HNRNPA2B1 (human) mapping to 7p15.2; Hnrnpa2b1 (mouse) mapping to 6 B3.

## SOURCE

hnRNP A2/B1 (C-3) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 220-239 within an internal region of hnRNP A2/B1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-393674 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

hnRNP A2/B1 (C-3) is recommended for detection of hnRNP A2/B1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for hnRNP A2/B1 siRNA (h): sc-43841, hnRNP A2/B1 siRNA (m): sc-43842, hnRNP A2/B1 shRNA Plasmid (h): sc-43841-SH, hnRNP A2/B1 shRNA Plasmid (m): sc-43842-SH, hnRNP A2/B1 shRNA (h) Lentiviral Particles: sc-43841-V and hnRNP A2/B1 shRNA (m) Lentiviral Particles: sc-43842-V.


Molecular Weight of hnRNP A2/B1: 36/38 kDa.

Positive Controls: K-562 nuclear extract: sc-2130, THP-1 nuclear extract: sc-24963 or HeLa nuclear extract: sc-2120.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



hnRNP A2/B1 (C-3): sc-393674. Western blot analysis of hnRNP A2/B1 expression in HeLa nuclear extract (A) and MEG-01 (B), Caco-2 (C), NIH/3T3 (D), BW5147 (E) and L6 (F) whole cell lysates.



hnRNP A2/B1 (C-3): sc-393674. Western blot analysis of hnRNP A2/B1 expression in A549 whole cell lysate (A) and HL-60 (B), MCF7 (C), Hep G2 (D) and Jurkat (E) nuclear extracts.

## SELECT PRODUCT CITATIONS

- Caggiano, C., et al. 2019. c-MYC empowers transcription and productive splicing of the oncogenic splicing factor Sam68 in cancer. *Nucleic Acids Res.* 47: 6160-6171.



See **hnRNP A2/B1 (B-7): sc-374053** for hnRNP A2/B1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.