

MTMR3 (A-12): sc-393779

BACKGROUND

Myotubularin and the myotubularin-related proteins (MTMR1-9) belong to a highly conserved family of eukaryotic phosphatases. They are protein tyrosine phosphatases that utilize inositol phospholipids, rather than phosphoproteins, as substrates. MTMR family members hydrolyze both phosphatidylinositol 3-phosphate (PtdIns3P) and PtdIns(3,5)P₂. MTMR2 interacts with MTMR5, an inactive family member that increases the enzymatic activity of MTMR2 and dictates its subcellular localization. Mutations in MTMR2 cause autosomal recessive Charcot-Marie-Tooth type 4B1 (CMT4B1), which is characterized by reduced nerve conduction velocities, focally folded myelin sheaths and demyelination. MTMR3 and MTMR4 can either interact with each other or self associate. MTMR6 regulates the activity of the calcium-activated potassium channel 3.1. MTMR9 regulates the activity of MTMR7 and MTMR8.

REFERENCES

1. Laporte, J., et al. 1997. Mutations in the MTM1 gene implicated in X-linked myotubular myopathy. ENMC International Consortium on Myotubular Myopathy. European Neuro-Muscular Center. Hum. Mol. Genet. 6: 1505-1511.
2. Blondeau, F., et al. 2000. Myotubularin, a phosphatase deficient in myotubular myopathy, acts on phosphatidylinositol 3-kinase and phosphatidylinositol 3-phosphate pathway. Hum. Mol. Genet. 9: 2223-2229.
3. Kim, S.A., et al. 2003. Regulation of myotubularin-related (MTMR)2 phosphatidylinositol phosphatase by MTMR5, a catalytically inactive phosphatase. Proc. Natl. Acad. Sci. USA 100: 4492-4497.
4. Mochizuki, Y. and Majerus, P.W. 2003. Characterization of myotubularin-related protein 7 and its binding partner, myotubularin-related protein 9. Proc. Natl. Acad. Sci. USA 100: 9768-9773.

CHROMOSOMAL LOCATION

Genetic locus: MTMR3 (human) mapping to 22q12.2; Mtmr3 (mouse) mapping to 11 A1.

SOURCE

MTMR3 (A-12) is a mouse monoclonal antibody raised against amino acids 6-265 mapping near the N-terminus of MTMR3 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

MTMR3 (A-12) is available conjugated to agarose (sc-393779 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-393779 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-393779 PE), fluorescein (sc-393779 FITC), Alexa Fluor® 488 (sc-393779 AF488), Alexa Fluor® 546 (sc-393779 AF546), Alexa Fluor® 594 (sc-393779 AF594) or Alexa Fluor® 647 (sc-393779 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-393779 AF680) or Alexa Fluor® 790 (sc-393779 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

MTMR3 (A-12) is recommended for detection of MTMR3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MTMR3 siRNA (h): sc-61088, MTMR3 siRNA (m): sc-61089, MTMR3 shRNA Plasmid (h): sc-61088-SH, MTMR3 shRNA Plasmid (m): sc-61089-SH, MTMR3 shRNA (h) Lentiviral Particles: sc-61088-V and MTMR3 shRNA (m) Lentiviral Particles: sc-61089-V.


Molecular Weight of MTMR3: 134 kDa.

Positive Controls: SK-N-MC cell lysate: sc-2237, human skeletal muscle extract: sc-363776 or mouse brain extract: sc-2253.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



MTMR3 (A-12): sc-393779. Western blot analysis of MTMR3 expression in SK-N-MC whole cell lysate (A) and human skeletal muscle tissue extract (B).



MTMR3 (A-12): sc-393779. Western blot analysis of MTMR3 expression in Sol8 nuclear extract (A) and mouse brain tissue extract (B).

SELECT PRODUCT CITATIONS

1. Jiang, Z., et al. 2023. Distinct shared and compartment-enriched oncogenic networks drive primary versus metastatic breast cancer. Nat. Commun. 14: 4313.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.