

Pontin 52 (A-4): sc-393854

BACKGROUND

Pontin 52 is nuclear matrix protein that is primarily expressed in the nucleus and is also present in the cytoplasm. Pontin 52 is expressed in the nucleoplasm of whole cells, but is not present in the nucleoli. Pontin 52 is also designated RUVBL1, for *E. coli* RuvB-like 1 protein, or NMP 238, and is the human homolog of rat TIP49. Pontin 52 contains an ATPase/helicase motif and may represent a class of cofactors recruited by transcriptional activation domains that function in diverse pathways. For instance, *in vivo*, Pontin 52 is complexed with Myc and Reptin 52, which is a Pontin 52 related protein also designated RUVBL2. The interaction of Pontin 52 with Myc is dependent upon a Myc domain essential for oncogenic activity, suggesting that functional Pontin 52 is an essential mediator of Myc oncogenic transformation. The gene encoding human Pontin 52 maps to chromosome 3q21.3.

REFERENCES

1. Bauer, A., et al. 1998. Pontin 52, an interaction partner of β -catenin, binds to the TATA box-binding protein. *Proc. Natl. Acad. Sci. USA* 95: 14787-14792.
2. Makino, Y., et al. 1998. TIP49, homologous to the bacterial DNA helicase RuvB, acts as an autoantigen in human. *Biochem. Biophys. Res. Commun.* 245: 819-823.
3. Holzmann, K., et al. 1998. Identification and characterization of the ubiquitously occurring nuclear matrix protein NMP 238. *Biochem. Biophys. Res. Commun.* 252: 39-45.
4. Qiu, X.B., et al. 1998. An eukaryotic RuvB-like protein (RUVBL1) essential for growth. *J. Biol. Chem.* 273: 27786-27793.
5. Lim, C.R., et al. 2000. The *Saccharomyces cerevisiae* RuvB-like protein, Tih2p, is required for cell cycle progression and RNA polymerase II-directed transcription. *J. Biol. Chem.* 275: 22409-22417.
6. Wood, M.A., et al. 2000. An ATPase/helicase complex is an essential cofactor for oncogenic transformation by c-Myc. *Mol. Cell* 5: 321-330.

CHROMOSOMAL LOCATION

Genetic locus: RUVBL1 (human) mapping to 3q21.3; Ruvbl1 (mouse) mapping to 6 D1.

SOURCE

Pontin 52 (A-4) is a mouse monoclonal antibody raised against amino acids 191-290 mapping within an internal region of Pontin 52 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-393854 X, 200 μ g/0.1 ml.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Pontin 52 (A-4) is recommended for detection of Pontin 52 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Pontin 52 (A-4) is also recommended for detection of Pontin 52 in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for Pontin 52 siRNA (h): sc-43543, Pontin 52 siRNA (m): sc-60010, Pontin 52 shRNA Plasmid (h): sc-43543-SH, Pontin 52 shRNA Plasmid (m): sc-60010-SH, Pontin 52 shRNA (h) Lentiviral Particles: sc-43543-V and Pontin 52 shRNA (m) Lentiviral Particles: sc-60010-V.

Pontin 52 (A-4) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Pontin 52: 50 kDa.

Positive Controls: SK-BR-3 cell lysate: sc-2218, K-562 nuclear extract: sc-2130 or KNRK whole cell lysate: sc-2214.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Pontin 52 (A-4): sc-393854. Western blot analysis of Pontin 52 expression in Jurkat (A), HT-1080 (B), BW5147 (C), RAT2 (D) and H19-7/IGF-IR (E) whole cell lysates and NIH/3T3 nuclear extract (F).



Pontin 52 (A-4): sc-393854. Western blot analysis of Pontin 52 expression in SK-BR-3 (A), NCI-H1299 (B) and KNRK (C) whole cell lysates and A-431 (D) and K-562 (E) nuclear extracts.

RESEARCH USE

For research use only, not for use in diagnostic procedures.