

caspase-10 p10 (E-11): sc-393983

BACKGROUND

A unique family of cysteine proteases has been described that differs in sequence, structure and substrate specificity from any previously described protease family. This family, CED-3/caspase-1, is comprised of caspase-1, caspase-2, caspase-3, caspase-4, caspase-6, caspase-7 (also designated Mch3, ICE-LAP3 or CMH-1), caspase-9 and caspase-10. CED-3/caspase-1 family members function as key components of the apoptotic machinery and act to destroy specific target proteins which are critical to cellular longevity. Poly(ADP-ribose) polymerase plays an integral role in surveying for DNA mutations and double strand breaks. Caspase-3, caspase-7 and caspase-9, but not caspase-1, cleave the nuclear protein PARP into an apoptotic fragment. Caspase-6, but not caspase-3, cleaves the nuclear lamins which are critical to maintaining the integrity of the nuclear envelope and cellular morphology. Caspase-10 activates caspase-3 and caspase-7 in response to apoptotic stimuli.

REFERENCES

1. Lindahl, T., et al. 1995. Post-translational modification of poly (ADP-ribose) polymerase induced by DNA strand breaks. *Trends Biochem. Sci.* 20: 405-411.
2. Duan, H., et al. 1996. ICE-LAP3, a novel mammalian homologue of the *Caenorhabditis elegans* cell death protein CED-3 is activated during FAS- and tumor necrosis factor-induced apoptosis. *J. Biol. Chem.* 271: 1621-1625.
3. Fernandes-Alnemri, T.F., et al. 1996. *In vitro* activation of CPP32 and Mch3 by Mch4, a novel human apoptotic cysteine protease containing two FADD-like domains. *Proc. Natl. Acad. Sci. USA* 93: 7464-7469.
4. Duan, H., et al. 1996. ICE-LAP6, a novel member of the ICE/CED-3 gene family, is activated by the cytotoxic T cell protease granzyme B. *J. Biol. Chem.* 271: 16720-16724.

CHROMOSOMAL LOCATION

Genetic locus: CASP10 (human) mapping to 2q33.1.

SOURCE

caspase-10 p10 (E-11) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 458-479 at the C-terminus of caspase-10 p10 of human origin.

PRODUCT

Each vial contains 200 µg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-393983 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

caspase-10 p10 (E-11) is recommended for detection of caspase-10 p10 subunit and precursor of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for caspase-10 siRNA (h): sc-29923, caspase-10 shRNA Plasmid (h): sc-29923-SH and caspase-10 shRNA (h) Lentiviral Particles: sc-29923-V.


Molecular Weight of caspase-10 p10: 58/10 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, Hep G2 cell lysate: sc-2227 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohisto-mount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



caspase-10 p10 (E-11): sc-393983. Western blot analysis of procaspase-10 expression in Jurkat (A), Hep G2 (B), HeLa (C), A-431 (D), K-562 (E) and NCI-H929 (F) whole cell lysates.



caspase-10 p10 (E-11): sc-393983. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing cytoplasmic staining of glandular cells (B).

SELECT PRODUCT CITATIONS

1. Cho, H.D., et al. 2019. Auricularin sensitizes primary prostate cancer cells to TRAIL-mediated apoptosis through up-regulation of the DR5-dependent pathway. *Food Chem. Toxicol.* 126: 223-232.

RESEARCH USE

For research use only, not for use in diagnostic procedures.