



IFN- ω siRNA (h): sc-39609

BACKGROUND

The genes encoding type I interferons (IFNs), which include 14 IFN- α genes, one IFN- β gene, one IFN- ω (also known as IFN- α II1) gene, and a number of IFN- ω pseudogenes, are clustered on human chromosome 9. Interferons- α and - β are cytokines that are widely known to induce potent anti-viral activity. IFN- α and - β exert a variety of other biological effects, including anti-tumor and immunomodulatory activities and are increasingly used clinically to treat a range of malignancies, myelodysplasias and autoimmune diseases. IFN- ω is antigenically different from human IFN- α , IFN- β or IFN- γ , but is a component of natural mixtures of IFN species produced by virus-induced leukocytes or Burkitt's lymphoma cells. The type I interferon receptor (IFN- α R) interacts with IFN- α , IFN- β and IFN- ω , and seems to be a multisubunit receptor.

REFERENCES

1. Adolf, G.R. 1987. Antigenic structure of human interferon ω 1 (interferon- α II1): comparison with other human interferons. *J. Gen. Virol.* 68: 1669-1676.
2. Lim, J.K., et al. 1994. Intrinsic ligand binding properties of the human and bovine α -interferon receptors. *FEBS Lett.* 350: 281-286.
3. Hussain, M., et al. 1996. Identification of interferon- α 7, - α 14 and - α 21 variants in the genome of a large human population. *J. Interferon Cytokine Res.* 16: 853-859.
4. Mire-Sluis, A.R., et al. 1996. An anti-cytokine bioactivity assay for interferons- α , - β and - ω . *J. Immunol. Methods* 195: 55-61.
5. Cutrone, E.C., et al. 1997. Contributions of cloned type I interferon receptor subunits to differential ligand binding. *FEBS Lett.* 404: 197-202.
6. Rozera, C., et al. 1999. Interferon (IFN)- β gene transfer into TS/A adenocarcinoma cells and comparison with IFN- α : differential effects on tumorigenicity and host response. *Am. J. Pathol.* 154: 1211-1222.
7. Barthe, C., et al. 2001. Expression of interferon- α (IFN- α) receptor 2c at diagnosis is associated with cytogenetic response in IFN- α -treated chronic myeloid leukemia. *Blood* 97: 3568-3573.
8. Eriksen, K.W., et al. 2004. Bi-phasic effect of interferon (IFN)- α : IFN- α up- and down-regulates interleukin-4 signaling in human T cells. *J. Biol. Chem.* 279: 169-176.
9. Suyama, T., et al. 2005. Upregulation of the interferon γ (IFN- γ)-inducible chemokines IFN-inducible T cell chemoattractant and monokine induced by IFN- γ and of their receptor CXCR3 in human renal cell carcinoma. *Cancer* 103: 258-267.

CHROMOSOMAL LOCATION

Genetic locus: IFNW1 (human) mapping to 9p21.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

IFN- ω siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IFN- ω shRNA Plasmid (h): sc-39609-SH and IFN- ω shRNA (h) Lentiviral Particles: sc-39609-V as alternate gene silencing products.

For independent verification of IFN- ω (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-39609A, sc-39609B and sc-39609C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IFN- ω siRNA (h) is recommended for the inhibition of IFN- ω expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IFN- ω gene expression knockdown using RT-PCR Primer: IFN- ω (h)-PR: sc-39609-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.