

IL-2 siRNA (h): sc-39619

BACKGROUND

Lymphokines are a group of signaling molecules involved in communication between cells of the immune system. Lymphokines secreted by activated lymphocytes include proteins such as interleukin-2. This protein is secreted primarily by helper T cells that have been activated through the T cell receptor complex or by other mitogens. Cells targeted by IL-2 include activated T helper and cytotoxic T cells, inducing their proliferation. The secretion of IL-2 can also act as a growth factor for B cells. To date, three different IL-2-dependent signal transduction pathways have been identified: the c-Fos/c-Jun induction pathway mediated by Src family protein-tyrosine kinases, the c-Myc induction pathway and the rapamycin-sensitive pathway, all of which result in the induction of Bcl-2. In addition, the transcription factor NFAT has been shown to play a major role in the regulation of IL-2 transcription and correlates to an age-related decline in the expression of IL-2.

REFERENCES

1. Smith, K.A. 1980. T cell growth factor. *Immunol. Rev.* 51: 337-357.
2. Taniguchi, T., et al. 1983. Structure and expression of a cloned cDNA for human interleukin-2. *Nature* 302: 305-310.
3. Lowenthal, J.W., et al. 1985. Similarities between interleukin-2 receptor number and affinity on activated B and T lymphocytes. *Nature* 315: 669-672.
4. Guy, G.R., et al. 1990. Lymphokine signal transduction. *Prog. Growth Factor Res.* 2: 45-70.

CHROMOSOMAL LOCATION

Genetic locus: IL2 (human) mapping to 4q27.

PRODUCT

IL-2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IL-2 shRNA Plasmid (h): sc-39619-SH and IL-2 shRNA (h) Lentiviral Particles: sc-39619-V as alternate gene silencing products.

For independent verification of IL-2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-39619A, sc-39619B and sc-39619C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IL-2 siRNA (h) is recommended for the inhibition of IL-2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

IL-2 (F-5): sc-133118 is recommended as a control antibody for monitoring of IL-2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG λ BP-HRP: sc-516132 or m-IgG λ BP-HRP (Cruz Marker): sc-516132-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG λ BP-FITC: sc-516185 or m-IgG λ BP-PE: sc-516186 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IL-2 gene expression knockdown using RT-PCR Primer: IL-2 (h)-PR: sc-39619-PR (20 μ l, 326 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Kim, S.D., et al. 2020. The malignancy of liver cancer cells is increased by IL-4/ERK/Akt signaling axis activity triggered by irradiated endothelial cells. *J. Radiat. Res.* 61: 376-387.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.