

IL-6 siRNA (m): sc-39628

BACKGROUND

Interleukin-6, or IL-6, is a multifunctional protein, 212 amino acids in length, that plays critical roles in host defense, immune response and hematopoiesis. IL-6 is constitutively expressed by epidermal Langerhans cells and its expression is induced in stimulated keratinocytes. IL-6, IL-1 β and TNF α act as endogenous pyrogens, regulating the fever response to bacterial invasion. The IL-6 receptor is a trimeric complex composed of an IL-6-specific α chain and a homodimer of the gp130 glycoprotein common to the IL-6, IL-11, CNTF, OSM and LIF receptors. Stimulation with IL-6 leads to gp130 homodimerization and the activation of associated kinases JAK1 and JAK2. Once activated, JAK1 and JAK2 phosphorylate Stat3, causing its nuclear translocation and transcription of Stat3-responsive genes. IL-6 has also been shown to activate the Ras/MAP kinase pathway, which regulates NFIL6 transcription.

REFERENCES

- Hirano, T., et al. 1986. Complementary DNA for a novel human interleukin (BSF-2) that induces B lymphocytes to produce immunoglobulin. *Nature* 324: 73-76.
- Nakajima, T., et al. 1993. Phosphorylation at Threonine 235 by a Ras-dependent mitogen-activated protein kinase cascade is essential for transcription factor NFIL6. *Proc. Natl. Acad. Sci. USA* 90: 2207-2211.

CHROMOSOMAL LOCATION

Genetic locus: Il6 (mouse) mapping to 5 B1.

PRODUCT

IL-6 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IL-6 shRNA Plasmid (m): sc-39628-SH and IL-6 shRNA (m) Lentiviral Particles: sc-39628-V as alternate gene silencing products.

For independent verification of IL-6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-39628A, sc-39628B and sc-39628C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IL-6 siRNA (m) is recommended for the inhibition of IL-6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

IL-6 (C12-1-hIL-6): sc-32296 is recommended as a control antibody for monitoring of IL-6 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IL-6 gene expression knockdown using RT-PCR Primer: IL-6 (m)-PR: sc-39628-PR (20 μ l, 542 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATION

- Tanimoto, T., et al. 2009. Genetic ablation of the Bach1 gene reduces hyperoxic lung injury in mice: role of IL-6. *Free Radic. Biol. Med.* 46: 1119-1126.
- Du, Y., et al. 2019. Activation of T lymphocytes as a novel mechanism in β_1 -adrenergic receptor autoantibody-induced cardiac remodeling. *Cardiovasc. Drugs Ther.* 33: 149-161.
- Masjedi, A., et al. 2020. Silencing of IL-6 and STAT3 by siRNA loaded hyaluronate-N,N-trimethyl chitosan nanoparticles potently reduces cancer cell progression. *Int. J. Biol. Macromol.* 149: 487-500.
- Salimifard, S., et al. 2020. Codelivery of BV6 and anti-IL6 siRNA by hyaluronate-conjugated PEG-chitosan-lactate nanoparticles inhibits tumor progression. *Life Sci.* 260: 118423.
- Allahyari, S.E., et al. 2021. Simultaneous inhibition of CD73 and IL-6 molecules by siRNA-loaded nanoparticles prevents the growth and spread of cancer. *Nanomedicine* 34: 102384.

RESEARCH USE

For research use only, not for use in diagnostic procedures.