

# IL-12B p40 siRNA (m): sc-39641

## BACKGROUND

The interleukins (ILs) are a broad family of well characterized cytokines, primarily of hematopoietic cell origin. ILs are secreted by immune cells (mainly macrophages, B cells or T cells) that regulate a wide range of immune system functions. The functions of different ILs vary from regulating inflammatory and immune responses, functioning as an autocrine factor and regulating and/or inhibiting other ILs. IL-12 is responsible for the differentiation of naive CD4<sup>+</sup> T cells into type 1 helper T cells that produce interferon- $\gamma$  (IFN- $\gamma$ ). It also activates production of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) from T and natural killer (NK) cells. IL-12 is a heterodimer composed of subunits IL-12A p35 and IL-12B p40. The p40 subunit of IL-12 also combines with p19, a protein that shows no biological activity by itself, to form a biologically active, composite cytokine, IL-23. IL-23 shares some *in vivo* functions with IL-12, including activation of the transcription factor Stat4 and IFN- $\gamma$  production and proliferation in PHA blast T cells, as well as in CD45RO (memory) T cells.

## REFERENCES

1. Oppmann, B., et al. 2000. Novel p19 protein engages IL-12 p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity* 13: 715-725.
2. Wiekowski, M.T., et al. 2001. Ubiquitous transgenic expression of the IL-23 subunit p19 induces multiorgan inflammation, runting, infertility, and premature death. *J. Immunol.* 166: 7563-7570.
3. Cooper, A.M., et al. 2002. Mice lacking bioactive IL-12 can generate protective, antigen-specific cellular responses to mycobacterial infection only if the IL-12 p40 subunit is present. *J. Immunol.* 168: 1322-1327.

## CHROMOSOMAL LOCATION

Genetic locus: IL12b (mouse) mapping to 11 B1.1.

## PRODUCT

IL-12B p40 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IL-12B p40 shRNA Plasmid (m): sc-39641-SH and IL-12B p40 shRNA (m) Lentiviral Particles: sc-39641-V as alternate gene silencing products.

For independent verification of IL-12B p40 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-39641A, sc-39641B and sc-39641C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

IL-12B p40 siRNA (m) is recommended for the inhibition of IL-12B p40 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

IL-12B p40 (F-10): sc-365389 is recommended as a control antibody for monitoring of IL-12B p40 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IL-12B p40 gene expression knockdown using RT-PCR Primer: IL-12B p40 (m)-PR: sc-39641-PR (20  $\mu$ l, 596 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Tan, M.S., et al. 2014. IL12/23 p40 inhibition ameliorates Alzheimer's disease-associated neuropathology and spatial memory in SAMP8 mice. *J. Alzheimers Dis.* 38: 633-646.
2. Sharma, N., et al. 2019. Deficiency of IL12p40 (interleukin 12 p40) promotes Ang II (angiotensin II)-induced abdominal aortic aneurysm. *Arterioscler. Thromb. Vasc. Biol.* 39: 212-223.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.