

4.1B (B-6): sc-398089



The Power to Question

BACKGROUND

The 4.1 gene family encodes a group of multifunctional cytoskeletal proteins (4.1R, 4.1G, 4.1N and 4.1B), which are predominantly expressed in the nervous system. 4.1G is a protein that stabilizes spectrin-Actin interactions and is associated with hereditary elliptocytosis. Red blood cell 4.1, designated 4.1R, is a multifunctional protein that is essential for maintaining erythrocyte shape and membrane mechanical properties. Both 4.1R and 4.1G are distributed in a unique pattern in the cerebellum and are believed to modulate the membrane mechanical properties of neuronal cells by promoting fodrin/Actin association. 4.1N and 4.1B, designated EPB41L1 and EPB41L3, respectively, are strongly expressed in the brain. Antibodies to 4.1N have been reported to detect multiple forms, each enriched in postsynaptic density preparations relative to brain homogenate. Antibodies to 4.1B have been reported to detect two forms.

REFERENCES

1. Peters, L.L., et al. 1998. Four paralogous protein 4.1 genes map to distinct chromosomes in mouse and human. *Genomics* 54: 348-350.
2. Takakuwa, Y. 2000. Protein 4.1, a multifunctional protein of the erythrocyte membrane skeleton: structure and functions in erythrocytes and nonerythroid cells. *Int. J. Hematol.* 72: 298-309.
3. Ohara, R., et al. 2000. Type II brain 4.1 (4.1B/KIAA0987), a member of the protein 4.1 family, is localized to neuronal paranodes. *Brain Res. Mol. Brain Res.* 85: 41-52.
4. Kontogianni-Konstantopoulos, A., et al. 2001. The prototypical 4.1R-10-kDa domain and the 4.1G-10-kDa paralog mediate fodrin-Actin complex formation. *J. Biol. Chem.* 276: 20679-20687.
5. Scott, C., et al. 2001. Protein 4.1 in forebrain postsynaptic density preparations: enrichment of 4.1 gene products and detection of 4.1R binding proteins. *Eur. J. Biochem.* 268: 1084-1094.

CHROMOSOMAL LOCATION

GGenic locus: EPB41L3 (human) mapping to 18p11.31; Epb41l3 (mouse) mapping to 17 E1.3.

SOURCE

4.1B (B-6) is a mouse monoclonal antibody raised against amino acids 654-810 mapping near the C-terminus of 4.1B of mouse origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

4.1B (B-6) is available conjugated to agarose (sc-398089 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398089 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398089 PE), fluorescein (sc-398089 FITC), Alexa Fluor® 488 (sc-398089 AF488), Alexa Fluor® 546 (sc-398089 AF546), Alexa Fluor® 594 (sc-398089 AF594) or Alexa Fluor® 647 (sc-398089 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398089 AF680) or Alexa Fluor® 790 (sc-398089 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

4.1B (B-6) is recommended for detection of 4.1B of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 4.1B siRNA (h): sc-40291, 4.1B siRNA (m): sc-40292, 4.1B shRNA Plasmid (h): sc-40291-SH, 4.1B shRNA Plasmid (m): sc-40292-SH, 4.1B shRNA (h) Lentiviral Particles: sc-40291-V and 4.1B shRNA (m) Lentiviral Particles: sc-40292-V.

Molecular Weight of 4.1B: 108 kDa.


Positive Controls: Neuro-2A whole cell lysate: sc-364185 or HEL 92.1.7 cell lysate: sc-2270.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



4.1B (B-6): sc-398089. Western blot analysis of 4.1B expression in Neuro-2A (A) and HEL 92.1.7 (B) whole cell lysates.



4.1B (B-6): sc-398089. Immunofluorescence staining of formalin-fixed SW480 cells showing membrane localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA