

YB-1 (E-7): sc-398146

BACKGROUND

Y-Box binding protein YB-1 (also known as CCAAT-binding transcription factor, enhancer factor I subunit A and DNA-binding protein B) belongs to a family of multifunctional proteins, which regulate both transcription and translation. Y-box proteins interact with a wide variety of nucleic acid structures to act as transcription factors and mRNA masking proteins. The modular structure of Y-box proteins includes a highly conserved N-terminal cold-shock domain (CSD, equivalent to the bacterial cold-shock proteins) and four basic C-terminal domains containing arginine clusters and aromatic residues. YB-1 plays a role in cell proliferation as an activator of growth-associated gene expression. YB-1 is also a repressor of the cell death-associated gene FAS. YB-1 may play an important role in controlling cell survival by regulating the expression of cell growth-associated and death-associated genes.

CHROMOSOMAL LOCATION

Genetic locus: YBX1 (human) mapping to 1p34.2; Ybx1 (mouse) mapping to 4 D2.1.

SOURCE

YB-1 (E-7) is a mouse monoclonal antibody raised against amino acids 196-240 mapping within an internal region of YB-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-398146 X, 200 µg/0.1 ml.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

YB-1 (E-7) is recommended for detection of YB-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for YB-1 siRNA (h): sc-38634, YB-1 siRNA (m): sc-38635, YB-1 siRNA (r): sc-63323, YB-1 shRNA Plasmid (h): sc-38634-SH, YB-1 shRNA Plasmid (m): sc-38635-SH, YB-1 shRNA Plasmid (r): sc-63323-SH, YB-1 shRNA (h) Lentiviral Particles: sc-38634-V, YB-1 shRNA (m) Lentiviral Particles: sc-38635-V and YB-1 shRNA (r) Lentiviral Particles: sc-63323-V.

YB-1 (E-7) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight (predicted) of YB-1: 36 kDa.


Molecular Weight (observed) of YB-1: 35-50 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, HeLa whole cell lysate: sc-2200 or SW480 cell lysate: sc-2219.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



YB-1 (E-7): sc-398146. Western blot analysis of YB-1 expression in HeLa (A) and Jurkat (B) whole cell lysates.



YB-1 (E-7): sc-398146. Western blot analysis of YB-1 expression in MCF7 (A) and SW480 (B) whole cell lysates.

SELECT PRODUCT CITATIONS

- Xiao, Y.Z., et al. 2020. Reducing hypothalamic stem cell senescence protects against aging-associated physiological decline. *Cell Metab.* 31: 534-548.e5.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.