

IL-3 (G-11): sc-398210

BACKGROUND

Interleukin-3, or IL-3, is a pleiotropic cytokine that is primarily secreted by activated T lymphocytes and stimulates the proliferation and differentiation of hematopoietic cells. IL-3 not only supports growth of both pluripotent stem cells and the more differentiated committed progenitors, but it also stimulates the functional activity of some fully differentiated cells. IL-3 has also been shown to protect mast cells from undergoing apoptosis. IL-3 exerts its biological effects through a receptor which consists of a ligand-specific α subunit and a signal transducing β subunit common to the IL-3/IL-5/GM-CSF receptors. The carboxy terminus of the β subunit has been shown to be necessary for activation of the MAP kinase signaling pathway. Although the IL-3 receptor has no intrinsic kinase activity, stimulation with IL-3 leads to tyrosine phosphorylation of the JAK/Tyk 2 family member, JAK2, which in turn activates and causes nuclear translocation of Stat5a and Stat5b.

REFERENCES

1. Abrams, J.S. and Pearce, M.K. 1988. Development of rat anti-mouse interleukin 3 monoclonal antibodies which neutralize bioactivity *in vitro*. *J. Immunol.* 140: 131-137.
2. Cockayne, D.A., et al. 1991. Antisense RNA inhibition of hematopoietic growth factor production. *Growth Factors* 5: 171-181.
3. Abrams, J.S., et al. 1992. Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. *Immunol. Rev.* 127: 5-24.
4. Magnelli, L., et al. 1993. Apoptosis induction in 32D cells by IL-3 withdrawal is preceded by a drop in the intracellular calcium level. *Biochem. Biophys. Res. Commun.* 194: 1394-1397.

CHROMOSOMAL LOCATION

Genetic locus: Il3 (mouse) mapping to 11 B1.3.

SOURCE

IL-3 (G-11) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 137-156 at the C-terminus of IL-3 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

IL-3 (G-11) is available conjugated to agarose (sc-398210 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398210 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398210 PE), fluorescein (sc-398210 FITC), Alexa Fluor[®] 488 (sc-398210 AF488), Alexa Fluor[®] 546 (sc-398210 AF546), Alexa Fluor[®] 594 (sc-398210 AF594) or Alexa Fluor[®] 647 (sc-398210 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-398210 AF680) or Alexa Fluor[®] 790 (sc-398210 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-398210 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

IL-3 (G-11) is recommended for detection of IL-3 of mouse origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for IL-3 siRNA (m): sc-39622, IL-3 shRNA Plasmid (m): sc-39622-SH and IL-3 shRNA (m) Lentiviral Particles: sc-39622-V.

Molecular Weight of IL-3: 15 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



IL-3 (G-11): sc-398210. Western blot analysis of mouse recombinant IL-3.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA