

# NUBPL (G-11): sc-398217

## BACKGROUND

NUBPL (nucleotide-binding protein-like) is a 319 amino acid mitochondrial protein that belongs to the Mrp/NBP35 ATP-binding protein family and exists as two alternatively spliced isoforms. Required for the assembly of the mitochondrial membrane respiratory chain NADH dehydrogenase (complex I), NUBPL may deliver one or more Fe-S clusters to complex I subunits. With highest expression in liver and kidney, NUBPL is expressed at significant levels in small intestine and brain. Defects in NUBPL are a cause of mitochondrial complex I deficiency (MT-C1D), a disorder of the mitochondrial respiratory chain that causes a wide range of afflictions from lethal neonatal disease to adult-onset neurodegenerative disorders. Phenotypes include macrocephaly with progressive leukodystrophy, non-specific encephalopathy, cardiomyopathy, myopathy, liver disease, Leigh syndrome, Leber hereditary optic neuropathy and some forms of Parkinson disease.

## REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 1987. Johns Hopkins University, Baltimore, MD. MIM Number: 252010. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Larsson, N.G., et al. 1991. Leber's hereditary optic neuropathy and complex I deficiency in muscle. *Ann. Neurol.* 30: 701-708.
3. Robinson, B.H. 1998. Human complex I deficiency: clinical spectrum and involvement of oxygen free radicals in the pathogenicity of the defect. *Biochim. Biophys. Acta* 1364: 271-286.

## CHROMOSOMAL LOCATION

Genetic locus: NUBPL (human) mapping to 14q12; Nubpl (mouse) mapping to 12 C1.

## SOURCE

NUBPL (G-11) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 46-69 within an internal region of NUBPL of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

NUBPL (G-11) is available conjugated to agarose (sc-398217 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398217 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398217 PE), fluorescein (sc-398217 FITC), Alexa Fluor® 488 (sc-398217 AF488), Alexa Fluor® 546 (sc-398217 AF546), Alexa Fluor® 594 (sc-398217 AF594) or Alexa Fluor® 647 (sc-398217 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398217 AF680) or Alexa Fluor® 790 (sc-398217 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-398217 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

NUBPL (G-11) is recommended for detection of NUBPL of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NUBPL (G-11) is also recommended for detection of NUBPL in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for NUBPL siRNA (h): sc-92266, NUBPL siRNA (m): sc-150092, NUBPL shRNA Plasmid (h): sc-92266-SH, NUBPL shRNA Plasmid (m): sc-150092-SH, NUBPL shRNA (h) Lentiviral Particles: sc-92266-V and NUBPL shRNA (m) Lentiviral Particles: sc-150092-V.


Molecular Weight of NUBPL isoforms: 34/18 kDa.

Positive Controls: NUBPL (h): 293T Lysate: sc-113739, T-47D cell lysate: sc-2293 or Hep G2 cell lysate: sc-2227.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



NUBPL (G-11): sc-398217. Western blot analysis of NUBPL expression in non-transfected: sc-117752 (A) and human NUBPL transfected: sc-113739 (B) 293T whole cell lysates.



NUBPL (G-11): sc-398217. Western blot analysis of NUBPL expression in T-47D (A) and Hep G2 (B) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.