

HS6ST1 (C-9): sc-398231

BACKGROUND

Heparan sulfate structures, which are responsible for executing multiple biologic activities, are generated and regulated by heparan sulfate (HS) biosynthetic enzymes. HS6ST1 (heparan sulfate 6-O-sulfotransferase 1), also known as HS6ST, is a 411 amino acid single-pass type II membrane protein that exists as multiple alternatively spliced isoforms and belongs to the sulfotransferase 6 family. Expressed in fetal brain, HS6ST1 functions as a 6-O-sulfation enzyme that specifically catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to the N-sulfoglucosamine residue (GlcNS) HS. HS6ST1 is subject to post-translational N-glycosylation and is encoded by a functional gene on human chromosome 2 and a pseudo-gene on human chromosome 1 (known as LOC728969).

REFERENCES

1. Habuchi, H., et al. 1998. Molecular characterization and expression of heparan-sulfate 6-sulfotransferase. Complete cDNA cloning in human and partial cloning in Chinese hamster ovary cells. *J. Biol. Chem.* 273: 9208-9213.
2. Habuchi, H., et al. 2000. The occurrence of three isoforms of heparan sulfate 6-O-sulfotransferase having different specificities for hexuronic acid adjacent to the targeted N-sulfoglucosamine. *J. Biol. Chem.* 275: 2859-2868.
3. Habuchi, H., et al. 2003. Biosynthesis of heparan sulphate with diverse structures and functions: two alternatively spliced forms of human heparan sulphate 6-O-sulphotransferase-2 having different expression patterns and properties. *Biochem. J.* 371: 131-142.
4. Edavettal, S.C., et al. 2004. A conformational change in heparan sulfate 3-O-sulfotransferase-1 is induced by binding to heparan sulfate. *Biochemistry* 43: 4680-4688.
5. Online Mendelian Inheritance in Man, OMIM™. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 604846. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Backen, A.C., et al. 2007. Heparan sulphate synthetic and editing enzymes in ovarian cancer. *Br. J. Cancer* 96: 1544-1548.
7. Nagai, N., et al. 2007. Regulation of heparan sulfate 6-O-sulfation by β -secretase activity. *J. Biol. Chem.* 282: 14942-14951.

CHROMOSOMAL LOCATION

Genetic locus: HS6ST1 (human) mapping to 2q14.3; Hs6st1 (mouse) mapping to 1 B.

SOURCE

HS6ST1 (C-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 52-79 within an internal region of HS6ST1 of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-398231 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

HS6ST1 (C-9) is recommended for detection of HS6ST1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HS6ST1 siRNA (h): sc-94375, HS6ST1 siRNA (m): sc-146089, HS6ST1 shRNA Plasmid (h): sc-94375-SH, HS6ST1 shRNA Plasmid (m): sc-146089-SH, HS6ST1 shRNA (h) Lentiviral Particles: sc-94375-V and HS6ST1 shRNA (m) Lentiviral Particles: sc-146089-V.

Molecular Weight (predicted) of HS6ST1: 48 kDa.


Molecular Weight (observed) of HS6ST1: 43 kDa.

Positive Controls: human brain extract: sc-364375 or Y79 cell lysate: sc-2240.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



HS6ST1 (C-9): sc-398231. Western blot analysis of HS6ST1 expression in human brain tissue extract.



HS6ST1 (C-9): sc-398231. Western blot analysis of HS6ST1 expression in Y79 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.