

CKAP2 (B-12): sc-398286

BACKGROUND

CKAP2 (cytoskeleton associated protein 2) is localized to the cytoplasm of humans and is expressed in tissues, including thymus and testis. CKAP2 is also referred to as LB1, TMAP or se20-10, and is a 682 amino acid protein which is expressed as three isoforms. CKAP2 is utilized during mitosis and is involved in regulating functions of microtubules, cellular death and the cell cycle. Before mitosis, CKAP2 is expressed in the cell cycle between phases G₁ and S, and accumulates between phases S and G₂. During mitosis, when the anaphase promoting complex is activated, CKAP2 is degraded. The regulation of CKAP2 is essential for proper spindle functions and cytokinesis, and it is thought that CKAP2 function is mediated via phosphorylation and dephosphorylation. Upon activation of p53 by CKAP2, the G₁ phase of the cell cycle is stopped, leading to cell death and apoptosis. Gastric tumors contain excessive amounts of CKAP2, which can lead to unregulated spindle functions and may be involved in the development and progression of gastric cancer.

REFERENCES

1. Udina, I.G., et al. 2001. Evolutionarily-conserved gene CKAP2, located in region 13q14.3 of the human genome, is frequently rearranged in various tumors. *Genetika* 37: 120-123.
2. Rakhmanaliev, E.R., et al. 2002. The structure of the human oncogenesis-associated CKAP2 (LB1) gene. *Mol. Biol.* 36: 985-989.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611569. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: CKAP2 (human) mapping to 13q14.3; Ckap2 (mouse) mapping to 8 A2.

SOURCE

CKAP2 (B-12) is a mouse monoclonal antibody raised against amino acids 431-494 mapping within an internal region of CKAP2 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CKAP2 (B-12) is available conjugated to agarose (sc-398286 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398286 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398286 PE), fluorescein (sc-398286 FITC), Alexa Fluor® 488 (sc-398286 AF488), Alexa Fluor® 546 (sc-398286 AF546), Alexa Fluor® 594 (sc-398286 AF594) or Alexa Fluor® 647 (sc-398286 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398286 AF680) or Alexa Fluor® 790 (sc-398286 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CKAP2 (B-12) is recommended for detection of CKAP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CKAP2 siRNA (h): sc-105207, CKAP2 siRNA (m): sc-142352, CKAP2 shRNA Plasmid (h): sc-105207-SH, CKAP2 shRNA Plasmid (m): sc-142352-SH, CKAP2 shRNA (h) Lentiviral Particles: sc-105207-V and CKAP2 shRNA (m) Lentiviral Particles: sc-142352-V.


Molecular Weight of CKAP2: 75 kDa.

Positive Controls: NTERA-2 cl.D1 whole cell lysate: sc-364181, U-698-M whole cell lysate: sc-364799 or Ramos cell lysate: sc-2216.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CKAP2 (B-12): sc-398286. Western blot analysis of CKAP2 expression in NTERA-2 cl.D1 (A), U-698-M (B), Ramos (C) and NAMALWA (D) whole cell lysates.



CKAP2 (B-12): sc-398286. Western blot analysis of CKAP2 expression in HEL 92.1.7 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Hotta, T., et al. 2022. EML2-S constitutes a new class of proteins that recognizes and regulates the dynamics of tyrosinated microtubules. *Curr. Biol.* 32: 3898-3910.e14.
2. Guo, L., et al. 2023. CKAP2 promotes cervical cancer progression by modulating the tumor microenvironment via NFκB signaling. *Am. J. Cancer Res.* 13: 2376-2391.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.