

NR5A2 (E-9): sc-398349

BACKGROUND

NR5A2 (nuclear receptor subfamily 5, group A, member 2), also designated B1-binding factor (B1F or B1F2), CYP7A promoter-binding factor (CPF), fetoprotein- α 1 (AFP) transcription factor (FTF, FTZ-F1 or FTZ-F1 β) and liver receptor homolog 1 (LRH-1), is a pre-adipocyte-specific nuclear receptor that regulates expression of aromatase in adipose tissue. NR5A2 belongs to the fushi tarazu factor-1 subfamily of orphan nuclear receptors. NR5A2 transcripts are abundant in the human ovary and testis and are predominantly expressed in tissues of endodermal origin. NR5A2 is a positive transcription factor for ABCG5 and ABCG8 and regulates genes involved in sterol and bile acid secretion from liver and intestine. It induces cell proliferation through the concomitant induction of cyclin D1 and E1, an effect that is potentiated by its interaction with β -catenin.

REFERENCES

1. Luo, Y., et al. 2001. The orphan nuclear receptor LRH-1 potentiates the sterol-mediated induction of the human CETP gene by liver X receptor. *J. Biol. Chem.* 276: 24767-24773.
2. Clyne, C.D., et al. 2002. Liver receptor homologue-1 (LRH-1) regulates expression of aromatase in preadipocytes. *J. Biol. Chem.* 277: 20591-20597.
3. Bohan, A., et al. 2003. Tumor necrosis factor α -dependent up-regulation of Lrh-1 and Mrp3(ABcc3) reduces liver injury in obstructive cholestasis. *J. Biol. Chem.* 278: 36688-36698.
4. Cai, Y.N., et al. 2003. LRH-1/hB1F and HNF1 synergistically up-regulate hepatitis B virus gene transcription and DNA replication. *Cell Res.* 13: 451-458.
5. Pezzi, V., et al. 2004. Differential expression of steroidogenic factor-1/adrenal 4 binding protein and liver receptor homolog-1 (LRH-1)/fetoprotein transcription factor in the rat testis: LRH-1 as a potential regulator of testicular aromatase expression. *Endocrinology* 145: 2186-2196.
6. Clyne, C.D., et al. 2004. Regulation of aromatase expression by the nuclear receptor LRH-1 in adipose tissue. *Mol. Cell. Endocrinol.* 215: 39-44.
7. Freeman, L.A., et al. 2004. The orphan nuclear receptor LRH-1 activates the ABCG5/ABCG8 intergenic promoter. *J. Lipid Res.* 45: 1197-1206.
8. Botrugno, O.A., et al. 2004. Synergy between LRH-1 and β -catenin induces G₁ cyclin-mediated cell proliferation. *Mol. Cell* 15: 499-509.

CHROMOSOMAL LOCATION

Genetic locus: NR5A2 (human) mapping to 1q32.1; Nr5a2 (mouse) mapping to 1 E4.

SOURCE

NR5A2 (E-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-15 at the N-terminus of NR5A2 of mouse origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-398349 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

NR5A2 (E-9) is recommended for detection of all NR5A2 isoforms of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NR5A2 siRNA (h): sc-37897, NR5A2 siRNA (m): sc-37898, NR5A2 shRNA Plasmid (h): sc-37897-SH, NR5A2 shRNA Plasmid (m): sc-37898-SH, NR5A2 shRNA (h) Lentiviral Particles: sc-37897-V and NR5A2 shRNA (m) Lentiviral Particles: sc-37898-V.

Molecular Weight of NR5A2 isoforms: 61/56/42 kDa.


Positive Controls: mouse pancreas extract: sc-364244.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



NR5A2 (E-9): sc-398349. Western blot analysis of NR5A2 expression in mouse pancreas tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.