

HIP-55 (E-5): sc-398358

BACKGROUND

Drebrins (developmentally regulated brain proteins) are cytoplasmic proteins that bind F-Actin in the brain and are involved in cell migration, extension of neuronal processes and plasticity of dendrites. HIP-55 (HPK1-interacting protein of 55 kDa), also known as ABP1, SH3P7 or DBNL (drebrin-like), is a 430 amino acid cytoplasmic protein that belongs to the ABP1 family. HIP-55 binds to F-Actin but is not involved in Actin polymerization, capping or bundling. In addition to containing an ADF-H domain, HIP-55 also consists of a SH3 domain, which mediates interaction with SHANK2, SHANK3 and PRAM-1. HIP-55 acts as an Actin-binding adapter protein and as a common effector of antigen receptor-signaling pathways in leukocytes. As a key component of the immunological synapse, HIP-55 regulates T-cell activation by bridging TCRs and the Actin cytoskeleton to gene activation and endocytic processes. HIP-55 is degraded by caspases during apoptosis.

REFERENCES

- Chen, Y.R., et al. 2001. Caspase-mediated cleavage of Actin-binding and SH3-domain-containing proteins cortactin, HS1, and HIP-55 during apoptosis. *Biochem. Biophys. Res. Commun.* 288: 981-989.
- Kessels, M.M., et al. 2001. Mammalian Abp1, a signal-responsive F-Actin-binding protein, links the Actin cytoskeleton to endocytosis via the GTPase dynamin. *J. Cell Biol.* 153: 351-366.
- Mise-Omata, S., et al. 2003. Mammalian Actin binding protein 1 is essential for endocytosis but not lamellipodia formation: functional analysis by RNA interference. *Biochem. Biophys. Res. Commun.* 301: 704-710.
- Han, J., et al. 2003. The SH3 domain-containing adaptor HIP-55 mediates c-Jun N-terminal kinase activation in T cell receptor signaling. *J. Biol. Chem.* 278: 52195-52202.
- Le Bras, S., et al. 2004. Recruitment of the Actin-binding protein HIP-55 to the immunological synapse regulates T cell receptor signaling and endocytosis. *J. Biol. Chem.* 279: 15550-15560.
- Denis, F.M., et al. 2005. PRAM-1 potentiates arsenic trioxide-induced JNK activation. *J. Biol. Chem.* 280: 9043-9048.
- Han, J., et al. 2005. HIP-55 is important for T cell proliferation, cytokine production, and immune responses. *Mol. Cell. Biol.* 25: 6869-6878.
- Le Bras, S., et al. 2007. Abl-SH3 binding protein 2, 3BP2, interacts with CIN85 and HIP-55. *FEBS Lett.* 581: 967-974.
- Haecckel, A., et al. 2008. The Actin-binding protein Abp1 controls dendritic spine morphology and is important for spine head and synapse formation. *J. Neurosci.* 28: 10031-10044.

CHROMOSOMAL LOCATION

Genetic locus: DBNL (human) mapping to 7p13; Dbnl (mouse) mapping to 11 A1.

SOURCE

HIP-55 (E-5) is a mouse monoclonal antibody raised against amino acids 100-186 mapping near the N-terminus of HIP-55 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

HIP-55 (E-5) is recommended for detection of HIP-55 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HIP-55 siRNA (h): sc-75255, HIP-55 siRNA (m): sc-75256, HIP-55 shRNA Plasmid (h): sc-75255-SH, HIP-55 shRNA Plasmid (m): sc-75256-SH, HIP-55 shRNA (h) Lentiviral Particles: sc-75255-V and HIP-55 shRNA (m) Lentiviral Particles: sc-75256-V.

Molecular Weight of HIP-55: 55 kDa.


Positive Controls: HIP-55 (h): 293T Lysate: sc-371066, K-562 whole cell lysate: sc-2203 or Raji whole cell lysate: sc-364236.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



HIP-55 (E-5): sc-398358. Western blot analysis of HIP-55 expression in Raji (A), K-562 (B), F9 (C), IB4 (D), A-10 (E) and KNRK (F) whole cell lysates.



HIP-55 (E-5): sc-398358. Western blot analysis of HIP-55 expression in non-transfected 293T: sc-117752 (A), human HIP-55 transfected 293T: sc-371066 (B) and Raji (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.