

α_1 B-AR siRNA (h): sc-39860

BACKGROUND

α_1 B-adrenergic receptors couple to $G_{q/11}$ and induce neoplastic transformation in NIH/3T3 cell transfectants. α_1 B receptors (α_1 B-ARs) can form hetero-oligomers with α_1 A and α_1 D receptors. α_1 B-AR transcripts are abundant in heart, brain, and kidney.

REFERENCES

- Allen, L.F., Lefkowitz, R.J., Caron, M.G. and Cotecchia, S. 1991. G protein-coupled receptor genes as protooncogenes: constitutively activating mutation of the α_1 B receptor enhances mitogenesis and tumorigenicity. *Proc. Natl. Acad. Sci. USA* 88: 11354-11358.
- Stanasila, L., Perez, J.B., Vogel, H. and Cotecchia, S. 2003. Oligomerization of the α_1 A- and α_1 B-adrenergic receptor subtypes. Potential implications in receptor internalization. *J. Biol. Chem.* 278: 40239-40251.
- Hague, C., Uberti, M.A., Chen, Z., Hall, R.A. and Minneman, K.P. 2004. Cell surface expression of α_1 D-adrenergic receptors is controlled by heterodimerization with α_1 B-adrenergic receptors. *J. Biol. Chem.* 279: 15541-15549.
- Papay, R., Gaivin, R., McCune, D.F., Rorabaugh, B.R., Macklin, W.B., McGrath, J.C. and Perez, D.M. 2004. Mouse α_1 B-adrenergic receptor is expressed in neurons and NG2 oligodendrocytes. *J. Comp. Neurol.* 478: 1-10.
- Daly, C.J. and McGrath, I. 2005. The role of the α_1 B-adrenergic receptor in vascular structure and function. *Hypertension* 45: e20-e21.
- Stanasila, L., Abuin, L., Diviani, D. and Cotecchia, S. 2006. Ezrin directly interacts with the α_1 B-adrenergic receptor and plays a role in receptor recycling. *J. Biol. Chem.* 281: 4354-4363.
- Shi, T., Gaivin, R.J., McCune, D.F., Gupta, M. and Perez, D.M. 2007. Dominance of the α_1 B-adrenergic receptor and its subcellular localization in human and TRAMP prostate cancer cell lines. *J. Recept. Signal Transduct. Res.* 27: 27-45.
- Takahashi, K., Hossain, M., Ahmed, M., Bhuiyan, M.A., Ohnuki, T. and Nagatomo, T. 2007. Asp 125 and Thr 130 in transmembrane domain 3 are major sites of α_1 B-adrenergic receptor antagonist binding. *Biol. Pharm. Bull.* 30: 1891-1894.

CHROMOSOMAL LOCATION

Genetic locus: ADRA1B (human) mapping to 5q33.3.

PRODUCT

α_1 B-AR siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see α_1 B-AR shRNA Plasmid (h): sc-39860-SH and α_1 B-AR shRNA (h) Lentiviral Particles: sc-39860-V as alternate gene silencing products.

For independent verification of α_1 B-AR (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-39860A, sc-39860B and sc-39860C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

α_1 B-AR siRNA (h) is recommended for the inhibition of α_1 B-AR expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor α_1 B-AR gene expression knockdown using RT-PCR Primer: α_1 B-ARR (h)-PR: sc-39860-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.