

# TCF-3 (F-5): sc-398640

## BACKGROUND

The TCF/LEF family of transcription factors are activated by the Wnt-1 and Wingless pathways and are characterized by the presence of a conserved protein motif, the high mobility group (HMG) 1 box, which mediates DNA binding. The TCF (T cell factor) proteins are required during developmental pathways. TCF-1 is essential for lymphoid cell development, while two other members, TCF-3 and TCF-4, are implicated in the development of the central nervous system. The Wnt mediated signaling pathway induces cytosolic  $\beta$ -catenin binding to TCF proteins within the nucleus, leading to the enhanced expression of the Wnt target genes. The  $\beta$ -catenin-TCF complexes are negatively regulated by the adenomatous polyposis coli (APC) tumor suppressor protein, which phosphorylates  $\beta$ -catenin and, in turn, increases the degradation of cytosolic  $\beta$ -catenin to, thereby, inhibit the activity of TCF proteins. Mutations in the APC gene, which are commonly observed in colorectal carcinomas, disrupt this regulatory pathway and correlate with an accumulation of  $\beta$ -catenin and the increased activation of the TCF target genes.

## REFERENCES

1. Van de Wetering, M., et al. 1991. Identification and cloning of TCF-1, a T lymphocyte-specific transcription factor containing a sequence-specific HMG box. *EMBO J.* 10: 123-132.
2. Van de Wetering, M., et al. 1992. The human T cell transcription factor-1 gene. Structure, localization, and promoter characterization. *J. Biol. Chem.* 267: 8530-8536.
3. Verbeek, S., et al. 1995. An HMG-box-containing T-cell factor required for thymocyte differentiation. *Nature* 374: 70-74.
4. Morin, P.J., et al. 1997. Activation of  $\beta$ -catenin-TCF signaling in colon cancer by mutations in  $\beta$ -catenin or APC. *Science* 275: 1787-1790.
5. Dorsky, R.I., et al. 1998. Control of neural crest cell fate by the Wnt signalling pathway. *Nature* 396: 370-373.
6. Young, C.S., et al. 1998. Wnt-1 induces growth, cytosolic  $\beta$ -catenin, and TCF/LEF transcriptional activation in Rat-1 fibroblasts. *Mol. Cell. Biol.* 18: 2474-2485.

## CHROMOSOMAL LOCATION

Genetic locus: TCF7L1 (human) mapping to 2p11.2; Tcf7l1 (mouse) mapping to 6 C1.

## SOURCE

TCF-3 (F-5) is a mouse monoclonal antibody raised against amino acids 421-575 of TCF-3 of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-398640 X, 200  $\mu$ g/0.1 ml.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

TCF-3 (F-5) is recommended for detection of TCF-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TCF-3 siRNA (h): sc-36618, TCF-3 siRNA (m): sc-36619, TCF-3 shRNA Plasmid (h): sc-36618-SH, TCF-3 shRNA Plasmid (m): sc-36619-SH, TCF-3 shRNA (h) Lentiviral Particles: sc-36618-V and TCF-3 shRNA (m) Lentiviral Particles: sc-36619-V.

TCF-3 (F-5) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TCF-3: 75 kDa.

Positive Controls: TCF-3 (h): 293T Lysate: sc-116647, MCF7 whole cell lysate: sc-2206 or 3T3-L1 cell lysate: sc-2243.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



TCF-3 (F-5): sc-398640. Western blot analysis of TCF-3 expression in MCF7 (A), 3T3-L1 (B) and Sol8 (C) whole cell lysates.



TCF-3 (F-5): sc-398640. Western blot analysis of TCF-3 expression in non-transfected: sc-117752 (A) and human TCF-3 transfected: sc-116647 (B) 293T whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.



See **TCF-3 (E-2): sc-166411** for TCF-3 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.