

DNA pol μ (E-8): sc-398666

BACKGROUND

DNA polymerase μ shares a number of characteristics with DNA polymerase β as well as with terminal deoxynucleotidyltransferase. Pol μ purportedly plays a role in microhomology mediated joining and the repair of double-stranded breaks. However, unlike other DNA polymerases, which show substrate specificity for deoxynucleotides, DNA Pol μ incorporates both deoxynucleotides and ribonucleotides in a template-directed manner. This unusual capability implies a novel role for this polymerase in DNA repair.

REFERENCES

- Chiu, A., et al. 2002. DNA polymerase μ gene expression in B-cell non-Hodgkin's lymphomas: an analysis utilizing *in situ* hybridization. *Am. J. Pathol.* 161: 1349-1355.
- Zhang, Y., et al. 2002. Lesion bypass activities of human DNA polymerase μ . *J. Biol. Chem.* 277: 44582-44587.
- Mahajan, K.N., et al. 2002. Association of DNA polymerase μ (pol μ) with Ku and ligase IV: role for pol μ in end-joining double-strand break repair. *Mol. Cell. Biol.* 22: 5194-5202.
- Nick McElhinny, S.A., et al. 2003. Polymerase μ is a DNA-directed DNA/RNA polymerase. *Mol. Cell. Biol.* 23: 2309-2315.
- Havener, J.M., et al. 2003. Translesion synthesis past platinum DNA adducts by human DNA polymerase μ . *Biochemistry* 42: 1777-1788.
- Ruiz, J.F., et al. 2004. Overexpression of human DNA polymerase μ (pol μ) in a Burkitt's lymphoma cell line affects the somatic hypermutation rate. *Nucleic Acids Res.* 32: 5861-5873.

CHROMOSOMAL LOCATION

Genetic locus: POLM (human) mapping to 7p13; Polm (mouse) mapping to 11 A1.

SOURCE

DNA pol μ (E-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 447-469 near the C-terminus of DNA polymerase μ of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

DNA pol μ (E-8) is available conjugated to agarose (sc-398666 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398666 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398666 PE), fluorescein (sc-398666 FITC), Alexa Fluor[®] 488 (sc-398666 AF488), Alexa Fluor[®] 546 (sc-398666 AF546), Alexa Fluor[®] 594 (sc-398666 AF594) or Alexa Fluor[®] 647 (sc-398666 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-398666 AF680) or Alexa Fluor[®] 790 (sc-398666 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-398666 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

DNA pol μ (E-8) is recommended for detection of DNA polymerase μ of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for DNA pol μ siRNA (h): sc-105304, DNA pol μ siRNA (m): sc-155886, DNA pol μ shRNA Plasmid (h): sc-105304-SH, DNA pol μ shRNA Plasmid (m): sc-155886-SH, DNA pol μ shRNA (h) Lentiviral Particles: sc-105304-V and DNA pol μ shRNA (m) Lentiviral Particles: sc-155886-V.

Positive Controls: A-431 nuclear extract: sc-2122, HeLa whole cell lysate: sc-2200 or JAR cell lysate: sc-2276.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



DNA pol μ (E-8): sc-398666. Western blot analysis of DNA pol μ expression in A-431 (A), Jurkat (B) and HeLa (C) nuclear extracts and Jurkat (D), HeLa (E) and JAR (F) whole cell lysates.



DNA pol μ (E-8): sc-398666. Western blot analysis of DNA pol μ expression in CCRF-CEM (A), Neuro-2A (B) and C6 (C) whole cell lysates.

SELECT PRODUCT CITATIONS

- Craxton, A., et al. 2018. PAXX and its paralogs synergistically direct DNA polymerase λ activity in DNA repair. *Nat. Commun.* 9: 3877.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA