

Rent3 (E-8): sc-398821

BACKGROUND

In eukaryotes, it is essential to have the ability to detect and degrade transcripts that lack full coding potential. Nonsense-mediated RNA decay (NMD) protects the organism by avoiding the translation of truncated peptides with dominant negative or deleterious gain-of-function potential. Rent1, a mammalian ortholog of Upf1, is essential for embryonic viability. Rent1 (also designated regulator of nonsense transcripts and HUpf1) contains an N-terminal zinc finger-like domain, NTPase domains and a region comprised of domains that define Rent1 as a superfamily group I helicase. Rent1 protein has nucleic-acid-dependent ATPase activity and 5' to 3' helicase activity. In addition, Rent1 is an RNA-binding protein whose activity is modulated by ATP and directly interacts with Rent2, which is a mammalian homolog of Upf2p. Two mammalian orthologs to Upf3p, Rent3a and Rent3b, are encoded by two separate genes. Rent3b (also known as Rent3X) is encoded by a X-linked gene and localizes primarily to the nucleus, while Rent 1 and Rent 2 localize primarily in the cytoplasm. Specific Rent3 protein interactions with Y14 and spliced mRNA suggest Rent3a and Rent3b serve as a link between splicing and NMD in the cytoplasm.

REFERENCES

1. Perlick, H.A., et al. 1996. Mammalian orthologues of a yeast regulator of nonsense transcript stability. *Proc. Natl. Acad. Sci. USA* 93: 10928-10932.
2. Page, M.F., et al. 1999. SMG-2 is a phosphorylated protein required for mRNA surveillance in *Caenorhabditis elegans* and related to Upf1p of yeast. *Mol. Cell. Biol.* 19: 5943-5951.

CHROMOSOMAL LOCATION

Genetic locus: UPF3A (human) mapping to 13q34, UPF3B (human) mapping to Xq24; Upf3a (mouse) mapping to 8 A1.1, Upf3b (mouse) mapping to X A3.3.

SOURCE

Rent3 (E-8) is a mouse monoclonal antibody raised against amino acids 51-240 mapping near the N-terminus of Rent3b of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-398821 X, 200 µg/0.1 ml.

Rent3 (E-8) is available conjugated to agarose (sc-398821 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398821 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398821 PE), fluorescein (sc-398821 FITC), Alexa Fluor® 488 (sc-398821 AF488), Alexa Fluor® 546 (sc-398821 AF546), Alexa Fluor® 594 (sc-398821 AF594) or Alexa Fluor® 647 (sc-398821 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398821 AF680) or Alexa Fluor® 790 (sc-398821 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Rent3 (E-8) is recommended for detection of Rent3a and Rent3b of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Rent3 (E-8) is also recommended for detection of Rent3a and Rent3b in additional species, including bovine.

Rent3 (E-8) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Rent3: 58 kDa.

Positive Controls: Rent3b (m): 293T Lysate: sc-123066, HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Rent3 (E-8): sc-398821. Western blot analysis of Rent3b expression in non-transfected 293T: sc-117752 (A), mouse Rent3b transfected 293T: sc-123066 (B), HeLa (C), Jurkat (D) and K-562 (E) whole cell lysates.



Rent3 (E-8): sc-398821. Western blot analysis of Rent3 expression in MOLT-4 (A), RAW 264.7 (B) and KNRK (C) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Rojo, C., et al. 2024. Caspases compromise SLU7 and UPF1 stability and NMD activity during hepatocarcinogenesis. *JHEP reports: innovation in hepatology* 6: 101118.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.