

# FHL-2 (E-8): sc-398866

## BACKGROUND

The four-and-a-half-LIM domain (FHL) proteins include FHL-1 (SLIM1), FHL-2 (SLIM3), FHL-3 (SLIM2) and FHL-4. The signature "half-domain", a single zinc finger domain located in the N-terminal region, differentiates FHLs from other LIM-only proteins, which have numbers of zinc fingers. Specific combinations of FHL proteins elicit selective activation of both CREB and CREM. Skeletal and cardiac muscle express FHL-1 in high levels as compared to the low level of expression in smooth muscle of the colon, small intestine and prostate. FHL-1 localizes to the cytosol of myoblasts, myotubes, and differentiated myocytes. FHL-2 is also located in cardiac and skeletal muscle, as well as in placenta and ovary tissues. FHL-3 is found in skeletal muscle, but absent in cardiac muscle. FHL-4 is expressed exclusively by the seminiferous epithelium of the testis, which suggests that FHL-4 is involved in spermatogenesis. The genetic loci for FHLs vary considerably despite similar amino acid sequences among the FHL group.

## REFERENCES

1. Morgan, M.J. and Madgwick, A.J. 1996. Slim defines a novel family of LIM-proteins expressed in skeletal muscle. *Biochem. Biophys. Res. Commun.* 225: 632-638.
2. Chan, K.K., et al. 1998. Molecular cloning and characterization of FHL-2, a novel LIM domain protein preferentially expressed in human heart. *Gene* 210: 345-350.
3. Lee, S.M., et al. 1998. Chromosomal mapping, tissue distribution and cDNA sequence of four-and-a-half LIM domain protein 1 (FHL1). *Gene* 216: 163-170.
4. Lee, S.M., et al. 1998. Chromosomal mapping of a skeletal muscle specific LIM-only protein FHL-3 to the distal end of the short arm of human chromosome 1. *Somat. Cell Mol. Genet.* 24: 197-202.
5. Morgan, M.J. and Madgwick, A.J. 1999. The LIM proteins FHL-1 and FHL-3 are expressed differently in skeletal muscle. *Biochem. Biophys. Res. Commun.* 255: 245-250.
6. Morgan, M.J. and Madgwick, A.J. 1999. The fourth member of the FHL family of LIM proteins is expressed exclusively in the testis. *Biochem. Biophys. Res. Commun.* 255: 251-255.
7. Greene, W.K., et al. 1999. Genomic structure, tissue expression and chromosomal location of the LIM-only gene, SLIM1. *Gene* 232: 203-207.

## CHROMOSOMAL LOCATION

Genetic locus: FHL2 (human) mapping to 2q12.1; Fhl2 (mouse) mapping to 1 B.

## SOURCE

FHL-2 (E-8) is a mouse monoclonal antibody raised against amino acids 1-57 mapping at the N-terminus of FHL-2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

FHL-2 (E-8) is recommended for detection of FHL-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FHL-2 siRNA (h): sc-37891, FHL-2 siRNA (m): sc-37892, FHL-2 shRNA Plasmid (h): sc-37891-SH, FHL-2 shRNA Plasmid (m): sc-37892-SH, FHL-2 shRNA (h) Lentiviral Particles: sc-37891-V and FHL-2 shRNA (m) Lentiviral Particles: sc-37892-V.


Molecular Weight of FHL-2: 32 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, human heart extract: sc-363763 or Sol8 cell lysate: sc-2249.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



FHL-2 (E-8): sc-398866. Western blot analysis of FHL-2 expression in T-47D (A), HT-1080 (B) and MG-63 (C) whole cell lysates and human heart tissue extract (D).



FHL-2 (E-8): sc-398866. Western blot analysis of FHL-2 expression in MCF7 (A), Sol8 (B) and L6 (C) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.