

# CSN7a (D-6): sc-398882

## BACKGROUND

The COP9 signalosome (CSN) complex is involved in several different developmental and cellular processes. The complex is made up of several widely expressed proteins: CSN1 (COPS1), CSN2 (COPS2), CSN3 (COPS3), CSN4 (COPS4), CSN5 (COPS5), CSN6 (COP6), CSN7a (COPS7, COPS7a), CSN7b (COP7b) and CSN8 (COP8). The CSN complex acts as a regulator for the ubiquitin conjugation pathway by mediating the deneddylation of the SCF-type E3 ligase complexes, which leads to a decrease in ubiquitin ligase activity of SCF-complexes. It is also involved in the phosphorylation of p53, c-Jun, I $\kappa$ B- $\alpha$  and IRF-8, as well as CSN-dependent phosphorylation of p53, and c-Jun protects and promotes degradation by the Ubl system. CSN7 is phosphorylated by CK2 and is composed of two subunits; a and b. CSN7a contains a PCI (proteasome CSN9 initiation factor 3) region, as well as a coiled-coil region and is predicted to interact with CSN2, CSN3, CSN4, CSN5, CSN6, CSN8, and GPS1. CSN7b contains only a PCI region and is predicted to interact with INT6.

## REFERENCES

- Seeger, M., et al. 1998. A novel protein complex involved in signal transduction possessing similarities to 26S Proteasome subunits. *FASEB J.* 12: 469-478.
- Bech-Otschir, D., et al. 2001. COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system. *EMBO J.* 20: 1630-1639.
- Hoareau Alves, K., et al. 2002. Association of the mammalian proto-oncoprotein Int-6 with the three protein complexes eIF3, COP9 signalosome and 26S Proteasome. *FEBS Lett.* 527: 15-21.

## CHROMOSOMAL LOCATION

Genetic locus: COPS7A (human) mapping to 12p13.31; Cops7a (mouse) mapping to 6 F2.

## SOURCE

CSN7a (D-6) is a mouse monoclonal antibody raised against amino acids 72-112 mapping within an internal region of CSN7a of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CSN7a (D-6) is available conjugated to agarose (sc-398882 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398882 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398882 PE), fluorescein (sc-398882 FITC), Alexa Fluor® 488 (sc-398882 AF488), Alexa Fluor® 546 (sc-398882 AF546), Alexa Fluor® 594 (sc-398882 AF594) or Alexa Fluor® 647 (sc-398882 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398882 AF680) or Alexa Fluor® 790 (sc-398882 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

CSN7a (D-6) is recommended for detection of CSN7a of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CSN7a siRNA (h): sc-60463, CSN7a siRNA (m): sc-60464, CSN7a shRNA Plasmid (h): sc-60463-SH, CSN7a shRNA Plasmid (m): sc-60464-SH, CSN7a shRNA (h) Lentiviral Particles: sc-60463-V and CSN7a shRNA (m) Lentiviral Particles: sc-60464-V.


Molecular Weight of CSN7a: 30 kDa.

Positive Controls: NTERA-2 cl.D1 whole cell lysate: sc-364181, HeLa whole cell lysate: sc-2200 or EOC 20 whole cell lysate: sc-364187.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



CSN7a (D-6): sc-398882. Western blot analysis of CSN7a expression in HeLa (A), NTERA-2 cl.D1 (B) and EOC 20 (C) whole cell lysates and human brain (D), rat placenta (E) and rat brain (F) tissue extracts.



CSN7a (D-6): sc-398882. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear and cytoplasmic localization (A). Immunofluorescence staining of formalin-fixed SW480 cells showing nuclear and cytoplasmic localization (B).

## SELECT PRODUCT CITATIONS

- Liu, R. and Moss, B. 2018. Vaccinia virus C9 ankyrin-repeat/F-box protein is a newly identified antagonist of the type I interferon-induced antiviral state. *J. Virol.* 92: e00053-18.
- Wang, J., et al. 2021. CSN7b defines a variant COP9 signalosome complex with distinct function in DNA damage response. *Cell Rep.* 34: 108662.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.