

# frizzled-3 siRNA (m): sc-39982

## BACKGROUND

The frizzled gene, originally identified in *Drosophila melanogaster*, is involved in the development of tissue polarity. The mammalian homolog of frizzled as well as several secreted mammalian frizzled-related proteins (FRPs) have been described. The frizzled proteins contain seven transmembrane domains, a cysteine-rich domain in the extracellular region and a carboxy terminal Ser/Thr-xxx-Val motif. They function as receptors for Wnt and are generally coupled to G proteins. The FRPs are involved in the Wnt signaling pathway by regulating the intracellular levels of  $\beta$ -catenin. Frizzled-3 (Fz-3) is a widely expressed, 666 amino acid protein, with relatively high expression in the central nervous system. Frizzled-3 has been associated with schizophrenia and has been shown to play a role in neural crest formation and hair follicle development. Two named isoforms of frizzled-3 exist as a result of alternative splicing events.

## REFERENCES

1. Kirikoshi, H., et al. 2000. Molecular cloning and genomic structure of human frizzled-3 at chromosome 8p21. *Biochem. Biophys. Res. Commun.* 271: 8-14.
2. Sala, C.F., et al. 2000. Identification, gene structure, and expression of human frizzled-3 (FZD3). *Biochem. Biophys. Res. Commun.* 273: 27-34.
3. Tan, C., et al. 2001. Kermit, a frizzled interacting protein, regulates frizzled-3 signaling in neural crest development. *Development* 128: 3665-3674.
4. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 606143. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Wang, Y., et al. 2002. frizzled-3 is required for the development of major fiber tracts in the rostral CNS. *J. Neurosci.* 22: 8563-8573.
6. Katsu, T., et al. 2003. The human frizzled-3 (FZD3) gene on chromosome 8p21, a receptor gene for Wnt ligands, is associated with the susceptibility to schizophrenia. *Neurosci. Lett.* 353: 53-56.
7. Jeong, S.H., et al. 2006. Investigation of genetic association between human frizzled homolog 3 gene (FZD3) and schizophrenia: results in a Korean population and evidence from meta-analysis. *Psychiatry Res.* 143: 1-11.

## CHROMOSOMAL LOCATION

Genetic locus: Fzd3 (mouse) mapping to 14 D1.

## PRODUCT

frizzled-3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see frizzled-3 shRNA Plasmid (m): sc-39982-SH and frizzled-3 shRNA (m) Lentiviral Particles: sc-39982-V as alternate gene silencing products.

For independent verification of frizzled-3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-39982A, sc-39982B and sc-39982C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

frizzled-3 siRNA (m) is recommended for the inhibition of frizzled-3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

frizzled-3 (C-1): sc-376105 is recommended as a control antibody for monitoring of frizzled-3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor frizzled-3 gene expression knockdown using RT-PCR Primer: frizzled-3 (m)-PR: sc-39982-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.