



EDAR siRNA (m): sc-40240

BACKGROUND

The tumor necrosis factor receptor (TNFR) superfamily represents a growing family of type I transmembrane glycoproteins that are involved in various cellular functions, including proliferation, differentiation and programmed cell death. These proteins share homology for cysteine-rich repeats in the extracellular ligand binding domain and the intracellular death domain. Members of the TNFR superfamily transmit signals through protein-protein interactions, and these signals can lead to the activation of either the caspase and Jun kinase pathways, which promote cell death, or the NF κ B pathway, which results in cell survival. The ectodermal dysplasia receptor (EDAR) promotes all three of these pathways and mediates ectodermal differentiation. EDAR is encoded by the downless gene and is mutated in ectodermal dysplasia syndromes, which are characterized by impaired hair, teeth and sweat gland development. Ectodysplasin A (EDA) is a type II membrane protein that is encoded by the Tabby gene and produces many splice variants, the longest of which, EDA-A1, serves as the ligand for EDAR. EDA-A2, which differs from EDA-A1 by the deletion of two amino acids, binds only the X-linked ectodysplasin-A2 receptor (XEDAR). Both EDAR and XEDAR exhibit homology with TROY.

REFERENCES

1. Gruss, H.J. 1996. Molecular, structural, and biological characteristics of the tumor necrosis factor ligand superfamily. *Int. J. Clin. Lab. Res.* 26: 143-159.
3. Gruss, H.J., et al. 1996. Structural and biological features of the TNF receptor and TNF ligand superfamilies: interactive signals in the pathobiology of Hodgkin's disease. *Ann. Oncol.* 7: 19-26.
4. Baker, S.J., et al. 1998. Modulation of life and death by the TNF receptor superfamily. *Oncogene* 17: 3261-3270.
5. Tucker, A.S., et al. 2000. EDAR/EDA interactions regulate enamel knot formation in tooth morphogenesis. *Development* 127: 4691-4700.
6. Yan, M., et al. 2000. Two-amino acid molecular switch in an epithelial morphogen that regulates binding to two distinct receptors. *Science* 290: 523-527.
7. Kojima, T., et al. 2000. TROY, a newly identified member of the tumor necrosis factor receptor superfamily, exhibits a homology with EDAR and is expressed in embryonic skin and hair follicles. *J. Biol. Chem.* 275: 20742-20747.
8. Kumar, A., et al. 2001. The ectodermal dysplasia receptor activates the nuclear factor- κ B, JNK, and cell death pathways and binds to ectodysplasin A. *J. Biol. Chem.* 276: 2668-2677.
9. Elomaa, O., et al. 2001. Ectodysplasin is released by proteolytic shedding and binds to the EDAR protein. *Hum. Mol. Genet.* 10: 953-962.

CHROMOSOMAL LOCATION

Genetic locus: EDAR (human) mapping to 2q11-q13; Edar (mouse) mapping to 10 B3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

EDAR siRNA (m) is a pool of 3 target-specific 20-25 nt siRNAs designed to knock down gene expression. Each vial contains 3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections.

For independent verification of EDAR (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3 nmol of lyophilized siRNA. These include: sc-40240A, sc-40240B and sc-40240C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

EDAR siRNA (m) is recommended for the inhibition of EDAR expression in mouse cells.

EDAR (C-20): sc-15290 is recommended as a control antibody for Western Blotting (starting dilution 1:100, dilution range 1:100-1:1,000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) protein detection using the recommended secondary reagents listed below.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 60 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238. Semi-quantitative RT-PCR may be performed using RT-PCR Primer: EDAR (m)-PR: sc-40240-PR (20 μ l).