



ACBP siRNA (m): sc-40311

BACKGROUND

Long chain acyl-CoA esters (LCAs) act as both substrates and intermediates in metabolism, and as regulators of various intracellular functions. Acyl-CoA binding protein (ACBP) specifically binds to LCA with high affinity and regulates its availability. ACBP is structurally and functionally conserved among a diverse group of organisms, including human, rat, frog, insect, plant and yeast. DBI, the gene encoding human ACBP, which maps to chromosome 2, is highly expressed in liver, soleus muscle and heart. The ACBP protein is also abundant in cells with a high level of lipogenesis and *de novo* fatty acid synthesis. Expression of ACBP is significantly induced during adipocyte differentiation. DBI is a target gene for proliferator-activated receptor (PPAR) γ , and is directly activated by PPAR γ /RXR α and PPAR α /RXR α , but not by PPAR δ /RXR α . In addition to acyl-CoA binding and transport, ACBP is also implicated in γ -aminobutyric acid type A receptor binding, steroidogenesis and peptide hormone release.

REFERENCES

- Knudsen, J. 1990. Acyl-CoA-binding protein (ACBP) and its relation to fatty acid-binding protein (FABP): an overview. *Mol. Cell. Biochem.* 98: 217-223.
- Knudsen, J., et al. 1993. The function of acyl-CoA-binding protein (ACBP)/diazepam binding inhibitor (DBI). *Mol. Cell. Biochem.* 123: 129-138.
- Gersuk, V.H., et al. 1995. Molecular cloning and chromosomal localization of a pseudogene related to the human acyl-CoA binding protein/diazepam binding inhibitor. *Genomics* 25: 469-476.
- Swinnen, J.V., et al. 1998. Identification of diazepam-binding Inhibitor/ Acyl-CoA-binding protein as a sterol regulatory element-binding protein-responsive gene. *J. Biol. Chem.* 273: 19938-19944.
- Knudsen, J., et al. 2000. Role of acyl-CoA binding protein in acyl-CoA metabolism and acyl-CoA-mediated cell signaling. *J. Nutr.* 130: 294S-298S.
- Helledie, T., et al. 2002. The gene encoding the Acyl-CoA-binding protein is activated by peroxisome proliferator-activated receptor γ through an intronic response element functionally conserved between humans and rodents. *J. Biol. Chem.* 277: 26821-26830.
- Franch, J., et al. 2002. Acyl-coenzyme A binding protein expression is fibre-type specific in rat skeletal muscle but not affected by moderate endurance training. *Pflugers Arch.* 443: 387-393.

CHROMOSOMAL LOCATION

Genetic locus: Dbi (mouse) mapping to 1 E2.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ACBP siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ACBP shRNA Plasmid (m): sc-40311-SH and ACBP shRNA (m) Lentiviral Particles: sc-40311-V as alternate gene silencing products.

For independent verification of ACBP (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-40311A, sc-40311B and sc-40311C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ACBP siRNA (m) is recommended for the inhibition of ACBP expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ACBP gene expression knockdown using RT-PCR Primer: ACBP (m)-PR: sc-40311-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.