

# $\gamma$ A-crystallin siRNA (m): sc-40451

## BACKGROUND

Crystallins, the major proteins of the vertebrate eye lens, are responsible for maintaining the transparency and the refractive index of the lens. Crystallins are divided into  $\alpha$ ,  $\beta$  and  $\gamma$  families, all of which usually contain seven distinctive protein regions, including four homologous motifs, one connecting peptide, and N- and C-terminal extensions. The  $\gamma$ -crystallin family is comprised of seven closely related proteins designated  $\gamma$ A-,  $\gamma$ B-,  $\gamma$ C-,  $\gamma$ D-,  $\gamma$ E-,  $\gamma$ F- and  $\gamma$ G-crystallin.  $\gamma$ A-crystallin, also known as CRYGA, CRYG5 or CRYG1, is a 174 amino acid member of the  $\gamma$ -crystallin family. Functioning as a monomer that has a two-domain  $\beta$  fold,  $\gamma$ A-crystallin, like other members of its family, plays a key role in ensuring the proper structure of the vertebrate eye lens. Defects in the gene encoding  $\gamma$ A-crystallin are associated with the formation of cataracts which are characterized by a clouding of the crystalline lens of the eye.

## REFERENCES

1. Brakenhoff, R.H., et al. 1990. Human  $\gamma$ -crystallin genes. A gene family on its way to extinction. *J. Mol. Biol.* 216: 519-532.
2. Hearne, C.M., et al. 1991. Trinucleotide repeat polymorphism at the CRYG1 locus. *Nucleic Acids Res.* 19: 5450.
3. Rogaev, E.I., et al. 1996. Linkage of polymorphic congenital cataract to the  $\gamma$ -crystallin gene locus on human chromosome 2q33-35. *Hum. Mol. Genet.* 5: 699-703.
4. Graw, J. 1997. The crystallins: genes, proteins and diseases. *Biol. Chem.* 378: 1331-1348.
5. Stöger, T., et al. 1997. The Cryner element in the murine  $\gamma$ -crystallin promoters interacts with lens proteins. *Ophthalmic Res.* 29: 161-171.
6. Slingsby, C. and Clout, N.J. 1999. Structure of the crystallins. *Eye* 13: 395-402.
7. Santhiya, S.T., et al. 2002. Novel mutations in the  $\gamma$ -crystallin genes cause autosomal dominant congenital cataracts. *J. Med. Genet.* 39: 352-358.
8. Salim, A. and Zaidi, Z.H. 2003. Homology models of human  $\gamma$ -crystallins: structural study of the extensive charge network in  $\gamma$ -crystallins. *Biochem. Biophys. Res. Commun.* 300: 624-630.
9. Messina-Baas, O.M., et al. 2006. Two affected siblings with nuclear cataract associated with a novel missense mutation in the CRYGD gene. *Mol. Vis.* 12: 995-1000.

## CHROMOSOMAL LOCATION

Genetic locus: Cryga (mouse) mapping to 1 C2.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

$\gamma$ A-crystallin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see  $\gamma$ A-crystallin shRNA Plasmid (m): sc-40451-SH and  $\gamma$ A-crystallin shRNA (m) Lentiviral Particles: sc-40451-V as alternate gene silencing products.

For independent verification of  $\gamma$ A-crystallin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-40451A, sc-40451B and sc-40451C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

$\gamma$ A-crystallin siRNA (m) is recommended for the inhibition of  $\gamma$ A-crystallin expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor  $\gamma$ A-crystallin gene expression knockdown using RT-PCR Primer:  $\gamma$ A-crystallin (m)-PR: sc-40451-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.