



DGAT1 siRNA (m): sc-40488

BACKGROUND

Glucose and Insulin are anabolic signals which upregulate the transcription of a series of lipogenic enzymes to convert excess carbohydrate into triglycerides for efficient energy storage. Acyl-Coenzyme A:diacylglycerol acyltransferase, also known as DGAT1 and ARGP1, is a microsomal enzyme that assists in the synthesis of fatty acids into triglycerides. DGAT catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol (DAG) and fatty acyl CoA as substrates. DGAT plays a fundamental role in the metabolism of cellular diacylglycerol and is important in higher eukaryotes for physiologic processes involving triacylglycerol metabolism such as intestinal fat absorption, lipoprotein assembly, adipose tissue formation and lactation. DGAT1 is involved in fat absorption in the intestine and in basal level triglyceride synthesis in adipose tissue, where it is more highly expressed. Mice lacking DGAT1 have increased energy expenditure and therefore are resistant to obesity. In addition, mice lacking both copies of DGAT1 are completely devoid of milk secretion, most likely because of deficient triglyceride synthesis in the mammary gland.

REFERENCES

1. Cases, S., et al. 1998. Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis. *Proc. Natl. Acad. Sci. USA* 95: 13018-13023.
2. Meegalla, R.L., et al. 2002. Concerted elevation of acyl-coenzyme A:diacylglycerol acyltransferase (DGAT) activity through independent stimulation of mRNA expression of DGAT1 and DGAT2 by carbohydrate and Insulin. *Biochem. Biophys. Res. Commun.* 298: 317-323.
3. Winter, A., et al. 2002. Association of a Lysine 232/alanine polymorphism in a bovine gene encoding acyl-CoA:diacylglycerol acyltransferase (DGAT1) with variation at a quantitative trait locus for milk fat content. *Proc. Natl. Acad. Sci. USA* 99: 9300-9305.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604900. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Chen, H.C., et al. 2003. Analysis of energy expenditure at different ambient temperatures in mice lacking DGAT1. *Am. J. Physiol. Endocrinol. Metab.* 284: E213-E218.

CHROMOSOMAL LOCATION

Genetic locus: Dgat1 (mouse) mapping to 15 D3.

PRODUCT

DGAT1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DGAT1 shRNA Plasmid (m): sc-40488-SH and DGAT1 shRNA (m) Lentiviral Particles: sc-40488-V as alternate gene silencing products.

For independent verification of DGAT1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-40488A, sc-40488B and sc-40488C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DGAT1 siRNA (m) is recommended for the inhibition of DGAT1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DGAT1 gene expression knockdown using RT-PCR Primer: DGAT1 (m)-PR: sc-40488-PR (20 μ l, 565 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Jung, H.J., et al. 2024. Inhibiting lipid droplet biogenesis enhances host protection against hypervirulent *Klebsiella pneumoniae* infections. *Med. Microbiol. Immunol.* 213: 26.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.