



PLC-L siRNA (m): sc-40846

BACKGROUND

Phosphoinositide-specific phospholipase C (PLC) plays a critical role in the initiation of receptor mediated signal transduction through the generation of the two second messengers, inositol 1, 4, 5-triphosphate and diacylglycerol from phosphatidylinositol 4, 5 bisphosphate. A total of nine mammalian PLC isozymes have been described. PLC-L (for PLC-deleted in lung carcinoma) is a PLC family member, the expression of which is decreased in small cell lung carcinomas. PLC-L is normally expressed in a variety of fetal and adult organs including the lung. PLC-L is thought to be involved in inositol phospholipid-based intracellular signaling cascade and aberrant expression of PLC-L may contribute to the genesis or progression of human lung carcinoma.

REFERENCES

1. Suh, P.G., et al. 1988. Inositol phospholipid-specific phospholipase C: complete cDNA and protein sequences and sequence homology to tyrosine kinase-related oncogene products. *Proc. Natl. Acad. Sci. USA* 85: 5419-5423.
2. Emori, Y., et al. 1989. A second type of rat phosphoinositide-specific phospholipase C containing asrc related sequence not essential for phosphoinositide-hydrolyzing activity. *J. Biol. Chem.* 264: 21885-21890.
3. Meldrum, E., et al. 1991. A second gene product of the inositol-phospholipid-specific phospholipase Cs subclass. *Eur. J. Biochem.* 196: 159-165.
4. Rhee, S.G. and Choi, K.D. 1992. Regulation of inositol phospholipid-specific phospholipase C isozymes. *J. Biol. Chem.* 267: 12393-12396.
5. Kim, M.J., et al. 1993. Cloning of cDNA encoding rat phospholipase C- β 4, a new member of the phospholipase C. *Biochem. Biophys. Res. Commun.* 194: 706-712.
6. Jhon, D., et al. 1993. Cloning, sequencing, purification and G_q -dependent activation of phospholipase C- β 3. *J. Biol. Chem.* 268: 6654-6661.
7. Kohno, T., et al. 1995. Identification of a novel phospholipase C family gene at chromosome 2q33 that is homozygously deleted in human small cell lung carcinoma. *Hum. Mol. Genet.* 4: 667-674.
8. Kelly, G.G., et al. 2001. Phospholipase C ϵ : a novel Ras effector. *EMBO J.* 20: 743-754.

CHROMOSOMAL LOCATION

Genetic locus: Plcl1 (mouse) mapping to 1 C1.2.

PRODUCT

PLC-L siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PLC-L shRNA Plasmid (m): sc-40846-SH and PLC-L shRNA (m) Lentiviral Particles: sc-40846-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PLC-L siRNA (m) is recommended for the inhibition of PLC-L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PLC-L gene expression knockdown using RT-PCR Primer: PLC-L (m)-PR: sc-40846-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.