

Polycystin-1 siRNA (h): sc-40861

BACKGROUND

Autosomal dominant polycystic kidney disease (ADPKD) is characterized by the formation of cysts in kidney tubules as well as in liver and pancreas tissues. Cells within these cysts display abnormalities in proliferation and polarity. The integral membrane protein, Polycystin-1 (PKD1) is mutated in a majority of patients with ADPKD. Polycystin-1 is expressed in renal tubular epithelial cells and colocalizes with cell and focal adhesion proteins, including E-cadherin, catenins, vinculin, and paxillin, to focal areas in order to form a larger multiprotein complex. Polycystin-1 is posttranslationally modified by tyrosine phosphorylation and associates with Polycystin-2 (PKD2) to mediate AP-1 expression, which suggests that Polycystin-1 is involved in cell-cell and cell-matrix interactions to control cell proliferation and polarity.

REFERENCES

1. Arnould, T., et al. 1998. The polycystic kidney disease 1 gene product mediates protein kinase C α -dependent and c-Jun N-terminal kinase-dependent activation of the transcription factor AP-1. *J. Biol. Chem.* 273: 6013-6018.
2. Huan, Y., et al. 1999. Polycystin-1, the PKD1 gene product, is in a complex containing E-cadherin and the catenins. *J. Clin. Invest.* 104: 1459-1468.
3. Ong, A.C., et al. 1999. Coordinate expression of the autosomal dominant polycystic kidney disease proteins, Polycystin-2 and Polycystin-1, in normal and cystic tissue. *Am. J. Pathol.* 154: 1721-1729.
4. Wilson, P.D., et al. 1999. The PKD1 gene product, "Polycystin-1," is a tyrosine-phosphorylated protein that colocalizes with α 2 β 1-Integrin in focal clusters in adherent renal epithelia. *Lab. Invest.* 79: 1311-1323.

CHROMOSOMAL LOCATION

Genetic locus: PKD1 (human) mapping to 16p13.3.

PRODUCT

Polycystin-1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Polycystin-1 shRNA Plasmid (h): sc-40861-SH and Polycystin-1 shRNA (h) Lentiviral Particles: sc-40861-V as alternate gene silencing products.

For independent verification of Polycystin-1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-40861A, sc-40861B and sc-40861C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Polycystin-1 siRNA (h) is recommended for the inhibition of Polycystin-1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Polycystin-1 (7E12): sc-130554 is recommended as a control antibody for monitoring of Polycystin-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Polycystin-1 gene expression knockdown using RT-PCR Primer: Polycystin-1 (h)-PR: sc-40861-PR (20 μ l, 375 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Hui, K., et al. 2014. The p38 MAPK-regulated PKD1/CREB/Bcl-2 pathway contributes to selenite-induced colorectal cancer cell apoptosis *in vitro* and *in vivo*. *Cancer Lett.* 354: 189-199.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.