



# Sck siRNA (h): sc-40928

## BACKGROUND

Src homology 2 (SH2) domains bind specifically to tyrosine-phosphorylated proteins that temporally participate in signal transduction events. Shc-like protein (Sck) is a neuronal adaptor protein that contains an N-terminal PTB (phosphotyrosine binding) domain, a collagen homology (CH) domain and a conserved C-terminal SH2 domain. Human Sck transcripts are present at high levels in liver, pancreas, prostate and ovary. In vascular endothelial cells, Sck participates in VEGF-induced signal transduction. Treatment of human umbilical vein endothelial (HUVEC) cells with VEGF induces recruitment of Sck to Tyrosine-1175 of the kinase insert domain-containing receptor (KDR) and enhances Sck tyrosine phosphorylation.

## REFERENCES

1. Kavanaugh, W.M. and Williams, L.T. 1994. An alternative to SH2 domains for binding tyrosine-phosphorylated proteins. *Science* 266: 1862-1865.
2. Nakamura, T., Muraoka, S., Sanokawa, R. and Mori, N. 1998. N-Shc and Sck, two neuronally expressed Shc adapter homologs: their differential regional expression in the brain and roles in neurotrophin and Src signaling. *J. Biol. Chem.* 273: 6960-6967.
3. Igarashi, K., Shigeta, K., Isohara, T., Yamano, T. and Uno, I. 1998. Sck interacts with KDR and Flt-1 via its SH2 domain. *Biochem. Biophys. Res. Commun.* 251: 77-82.
4. Warner, A.J., Lopez-Dee, J., Knight, E.L., Feramisco, J.R. and Prigent, S.A. 2000. The Shc-related adaptor protein, Sck, forms a complex with the vascular-endothelial-growth-factor receptor KDR in transfected cells. *Biochem. J.* 347: 501-509.
5. Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 605217. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Kojima, T., Yoshikawa, Y., Takada, S., Sato, M., Nakamura, T., Takahashi, N., Copeland, N.G., Gilbert, D.J., Jenkins, N.A. and Mori, N. 2001. Genomic organization of the Shc-related phosphotyrosine adapters and characterization of the full-length Sck/ShcB: specific association of p68-Sck/ShcB with pp135. *Biochem. Biophys. Res. Commun.* 284: 1039-1047.
7. Ratcliffe, K.E., Tao, Q., Yavuz, B., Stoletov, K.V., Spring, S.C. and Terman, B.I. 2002. Sck is expressed in endothelial cells and participates in vascular endothelial growth factor-induced signaling. *Oncogene* 21: 6307-6316.
8. LocusLink Report (LocusID: 25759). <http://www.ncbi.nlm.nih.gov/LocusLink/>

## CHROMOSOMAL LOCATION

Genetic locus: SHC2 (human) mapping to 19p13.3.

## PRODUCT

Sck siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Sck shRNA Plasmid (h): sc-40928-SH and Sck shRNA (h) Lentiviral Particles: sc-40928-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Sck siRNA (h) is recommended for the inhibition of Sck expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

Sck (E-3): sc-514627 is recommended as a control antibody for monitoring of Sck gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Sck gene expression knockdown using RT-PCR Primer: Sck (h)-PR: sc-40928-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.