

SOCS-7 siRNA (h): sc-41004

BACKGROUND

The SOCS (suppressor of cytokine signaling) gene family consists of a group of proteins that negatively regulate cytokine signal transduction. The SOCS family proteins contain a central SH2 domain and a carboxy-terminal region termed the "SOCS box." The SOCS-1 (also called SSI-1 and JAB), SOCS-2 (also called SSI-2 and CIS2) and SOCS-3 (also called SSI-3 and CIS3) genes are known to be upregulated by IL-6 and other cytokines. SOCS-4, SOCS-5, SOCS-6 and SOCS-7 were identified by their sequence homology with the SOCS box. CIS (cytokine-inducible SH2-containing protein) is also a member of the SOCS family.

REFERENCES

1. Yoshimura, A., et al. 1995. A novel cytokine-inducible gene CIS encodes an SH2-containing protein that binds to tyrosine-phosphorylated interleukin 3 and erythropoietin receptors. *EMBO J.* 14: 2816-2826.
2. Matsumoto, A., et al. 1997. CIS, a cytokine inducible SH2 protein, is a target of the JAK-Stat5 pathway and modulates Stat5 activation. *Blood* 89: 3148-3154.
3. Starr, R., et al. 1997. A family of cytokine-inducible inhibitors of signalling. *Nature* 387: 917-921.
4. Nicholson, S.E., et al. 1998. The SOCS proteins: a new family of negative regulators of signal transduction. *J. Leukoc. Biol.* 63: 665-668.
5. Hilton, D.J., et al. 1998. Twenty proteins containing a C-terminal SOCS box form five structural classes. *Proc. Natl. Acad. Sci. USA* 95: 114-119.

CHROMOSOMAL LOCATION

Genetic locus: SOCS7 (human) mapping to 17q12.

PRODUCT

SOCS-7 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SOCS-7 shRNA Plasmid (h): sc-41004-SH and SOCS-7 shRNA (h) Lentiviral Particles: sc-41004-V as alternate gene silencing products.

For independent verification of SOCS-7 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41004A, sc-41004B and sc-41004C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SOCS-7 siRNA (h) is recommended for the inhibition of SOCS-7 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

SOCS-7 (E-8): sc-137241 is recommended as a control antibody for monitoring of SOCS-7 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SOCS-7 gene expression knockdown using RT-PCR Primer: SOCS-7 (h)-PR: sc-41004-PR (20 μ l, 522 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.