

TPST-1 siRNA (m): sc-41076

BACKGROUND

The tyrosylprotein sulfotransferases TPST-1 and TPST-2 catalyze the sulfation of tyrosine residues within secreted and membrane-bound proteins, such as cell adhesion molecules, G protein-coupled receptors, coagulation factors, serpins, extracellular matrix proteins, and hormones. Although both TPST-1 and TPST-2 utilize 3'-phosphoadenosine 5'-phosphosulfate as their sulfate donor, they differ in their substrate specificity. The TPSTs are evolutionarily conserved proteins found in a wide variety of species, including human, mouse, *C. elegans*, and plants. They are ubiquitously expressed in several tissues, including liver, lung, heart, and cerebellum. Both TPST-1 and TPST-2 localize to the Golgi complex. Chronic alcohol consumption stimulates a threefold increase in TPST levels in the gastric mucosa and liver, indicating that TPST may play a role in alcoholism. The genes encoding human TPST-1 and TPST-2 map to chromosomes 7q11.21 and 22q12.1, respectively.

REFERENCES

1. Kasinathan, C., Sundaram, P., Slomiany, B.L. and Slomiany, A. 1993. Inhibition of tyrosylprotein sulfotransferase by sphingosine and its reversal by acidic phospholipids. *Biochemistry* 32: 1194-1198.
2. Ouyang, Y.B. and Moore, K.L. 1998. Molecular cloning and expression of human and mouse tyrosylprotein sulfotransferase-2 and a tyrosylprotein sulfotransferase homologue in *Caenorhabditis elegans*. *J. Biol. Chem.* 273: 24770-24774.
3. Beisswanger, R., Corbeil, D., Vannier, C., Thiele, C., Dohrmann, U., Kellner, R., Ashman, K., Niehrs, C. and Huttner, W.B. 1998. Existence of distinct tyrosylprotein sulfotransferase genes: molecular characterization of tyrosylprotein sulfotransferase-2. *Proc. Natl. Acad. Sci. USA* 95: 11134-11139.
4. Kasinathan, C., Ramaprasad, P., William, S. and Espina, N. 1998. Stimulation of tyrosylprotein sulfotransferase activity by ethanol: role of increased enzyme level. *Alcohol* 15: 271-276.
5. Hanai, H., Nakayama, D., Yang, H., Matsubayashi, Y., Hirota, Y. and Sakagami, Y. 2000. Existence of a plant tyrosylprotein sulfotransferase: novel plant enzyme catalyzing tyrosine O-sulfation of preprophytosulfokine variants *in vitro*. *FEBS Lett.* 470: 97-101.
6. Kehoe, J.W., Maly, D.J., Verdugo, D.E., Armstrong, J.I., Cook, B.N., Ouyang, Y.B., Moore, K.L., Ellman, J.A. and Bertozzi, C.R. 2002. Tyrosylprotein sulfotransferase inhibitors generated by combinatorial target-guided ligand assembly. *Bioorg. Med. Chem. Lett.* 12: 329-332.
7. Ouyang, Y.B., Crawley, J.T., Aston, C.E. and Moore, K.L. 2002. Reduced body weight and increased postimplantation fetal death in tyrosylprotein sulfotransferase-1-deficient mice. *J. Biol. Chem.* 277: 23781-23787.
8. LocusLink Report (LocusID: 8460). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: Tpst1 (mouse) mapping to 5 G1.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TPST-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TPST-1 shRNA Plasmid (m): sc-41076-SH and TPST-1 shRNA (m) Lentiviral Particles: sc-41076-V as alternate gene silencing products.

For independent verification of TPST-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41076A, sc-41076B and sc-41076C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TPST-1 siRNA (m) is recommended for the inhibition of TPST-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TPST-1 gene expression knockdown using RT-PCR Primer: TPST-1 (m)-PR: sc-41076-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.