

Wnt-3a siRNA (m): sc-41109

BACKGROUND

The Wnt gene family encodes secreted signaling molecules that bind to frizzled receptors and influence oncogenesis and developmental processes, including regulation of cell fate and patterning during embryogenesis. The Wnt family has two functional classes according to their biological activities; Wnts that signal through a Wnt-1/wingless pathway by stabilizing cytoplasmic β -catenin, and Wnts that stimulate intracellular Ca^{2+} release and activate two kinases, CamKII and PKC, in a G protein-dependent manner. Wnt-3a is an intercellular signaling molecule that mediates cytoskeletal reorganization and regulates hippocampal development. Human Wnt-3a is 96% homologous to mouse Wnt-3a protein and 84% homologous to human Wnt-3 protein. The human Wnt-3a gene clusters with the Wnt-14 gene at chromosome 1q42.13.

REFERENCES

1. Shibamoto, S., et al. 1998. Cytoskeletal reorganization by soluble Wnt-3a protein signalling. *Genes Cells* 3: 659-670.
2. Kuhl, M., et al. 2000. The Wnt/ Ca^{2+} pathway: a new vertebrate Wnt signaling pathway takes shape. *Trends Genet.* 16: 279-283.
3. Lee, S.M., et al. 2000. A local Wnt-3a signal is required for development of the mammalian hippocampus. *Development* 127: 457-467.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606359. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. LocusLink Report (LocusID: 89780). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: Wnt3a (mouse) mapping to 11 B1.3.

PRODUCT

Wnt-3a siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Wnt-3a shRNA Plasmid (m): sc-41109-SH and Wnt-3a shRNA (m) Lentiviral Particles: sc-41109-V as alternate gene silencing products.

For independent verification of Wnt-3a (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41109A, sc-41109B and sc-41109C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Wnt-3a siRNA (m) is recommended for the inhibition of Wnt-3a expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Wnt-3a (3A6): sc-136163 is recommended as a control antibody for monitoring of Wnt-3a gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Wnt-3a gene expression knockdown using RT-PCR Primer: Wnt-3a (m)-PR: sc-41109-PR (20 μl , 461 bp). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

SELECT PRODUCT CITATIONS

1. He, X., et al. 2016. TLR4 activation promotes bone marrow MSC proliferation and osteogenic differentiation via Wnt-3a and Wnt-5a signaling. *PLoS ONE* 11: e0149876.
2. Agarwal, S., et al. 2017. Fetuin-A downregulates adiponectin through Wnt-PPAR γ pathway in lipid induced inflamed adipocyte. *Biochim. Biophys. Acta Mol. Basis Dis.* 1863: 174-181.
3. Li, J., et al. 2022. Pulsed electromagnetic fields inhibit mandibular bone deterioration depending on the Wnt3a/ β -catenin signaling activation in type 2 diabetic db/db mice. *Sci. Rep.* 12: 7217.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.